arXiv:1302.6143v1 [math.NT] 25 Feb 2013

The Relation Between Global $\mathfrak{G}$-Shtukas and Local $\mathbb{P}$-Shtukas

Esmail Arasteh Rad and Urs Hartl

February 26, 2013

Abstract

In a sequence of two articles we propose to view the moduli stacks of global $\mathfrak{G}$-shtukas as function field analogs for Shimura varieties. Here $\mathfrak{G}$ is a parahoric Bruhat-Tits group scheme over a smooth projective curve, and global $\mathfrak{G}$-shtukas are generalizations of Drinfeld shtukas and analogs of abelian varieties with additional structure. Our moduli stacks generalize various moduli spaces used by different authors to prove instances of the Langlands program over function fields. In the present article we explain the relation between global $\mathfrak{G}$-shtukas and local $\mathbb{P}$-shtukas, which are the function field analogs of p-divisible groups with additional structure. We prove the analog of a theorem of Serre and Tate stating the equivalence between the deformations of a global $\mathfrak{G}$-shtuka and its associated local $\mathbb{P}$-shtukas. We explain the relation of local $\mathbb{P}$-shtukas with Galois representations through their Tate modules. And we prove the existence of Rapoport-Zink spaces for local $\mathbb{P}$-shtukas as formal schemes locally formally of finite type. In the sequel to this article we use the Rapoport-Zink spaces to uniformize the moduli stacks of global $\mathfrak{G}$-shtukas.

Mathematics Subject Classification (2000): 11G09, (11G18, 14L05, 14M15)

Contents

1 Introduction **2**

1.1 Notation and Conventions 3

2 Local $\mathbb{P}$-Shtukas and Global $\mathfrak{G}$-Shtukas **4**

2.1 Loop Groups and Local $\mathbb{P}$-Shtukas 4

2.2 Global $\mathfrak{G}$-Shtukas 7

3 Tate Modules for Local $\mathbb{P}$-Shtukas **8**

4 The Rapoport-Zink Spaces for Local $\mathbb{P}$-Shtukas **11**

4.1 Unbounded Rapoport-Zink Spaces 11

4.2 Bounded Local $\mathbb{P}$-Shtukas 14

4.3 Representability of the Bounded Rapoport-Zink Functor 18

5 The Relation Between Global $\mathfrak{G}$-Shtukas and Local $\mathbb{P}$-Shtukas **25**

5.1 Preliminaries on $\mathfrak{G}$-Torsors 25

5.2 The Global-Local Functor 27

5.3 The Analog of the Serre-Tate Theorem 30

References **31**

1 Introduction

Let $\mathbb{F}_q$ be a finite field with q elements, let C be a smooth projective geometrically irreducible curve over $\mathbb{F}_q$, and let $\mathfrak{G}$ be a parahoric Bruhat-Tits group scheme over C; see Definition 2.12. A *global* $\mathfrak{G}$-*shtuka* $\underline{\mathcal{G}}$ over an $\mathbb{F}_q$-scheme S is a tuple $(\mathcal{G}, s_1, \ldots, s_n, \tau)$ consisting of a $\mathfrak{G}$-torsor $\mathcal{G}$ over $C_S := C \times_{\mathbb{F}_q} S$, an n-tuple of (characteristic) sections $(s_1, \ldots, s_n) \in C^n(S)$ and a Frobenius connection τ defined outside the graph of the sections s_i's, that is, an isomorphism $\tau \colon \sigma^* \mathcal{G}|_{C_S \smallsetminus \cup_i \Gamma_{s_i}} \xrightarrow{\sim} \mathcal{G}|_{C_S \smallsetminus \cup_i \Gamma_{s_i}}$ where $\sigma^* = (\mathrm{id}_C \times \mathrm{Frob}_{q,S})^*$.

In [AH13] we will show that the moduli stack $\nabla_n \mathscr{H}^1(C, \mathfrak{G})$ of global $\mathfrak{G}$-shtukas, after imposing suitable boundedness conditions and level structures, is an algebraic Deligne-Mumford stack over C^n. Spelling out the Riemann-Hilbert correspondence for function fields, together with the tannakian philosophy, one sees that $\nabla_n \mathscr{H}^1(C, \mathfrak{G})$ may play the same role that Shimura varieties play for number fields. More specifically one can hope that the Langlands correspondence for function fields is realized on its cohomology. Note that in particular this moduli stack generalizes the space of F-sheaves $FSh_{D,r}$ which was considered by Drinfeld [Dri87] and Lafforgue [Laf02] in their proof of the Langlands correspondence for $\mathfrak{G} = \mathrm{GL}_2$ (resp. $\mathfrak{G} = \mathrm{GL}_r$), and which in turn was generalized by Varshavsky's [Var04] moduli stacks $FBun$. It likewise generalizes the moduli stacks $Cht_{\underline{\lambda}}$ of Ngô and Ngô Dac [NN08], $\mathcal{E}\ell\ell_{C,\mathscr{D},I}$ of Laumon, Rapoport and Stuhler [LRS93], and $\mathcal{A}b\text{-}\mathcal{S}h^{r,d}_H$ of the second author [Har05]; see [AH13, Remark 3.16]. On the other hand this analogy can be viewed as an attempt to build a bridge between the geometric Langlands program and the arithmetic Langlands program, where the role of global $\mathfrak{G}$-shtukas is played by abelian varieties (together with additional structures), respectively D-modules.

In [AH13] we also prove that $\nabla_n \mathscr{H}^1(C, \mathfrak{G})$ has a Rapoport-Zink uniformization by Rapoport-Zink spaces for local $\mathbb{P}$-shtukas, where $\mathbb{P}$ is a parahoric group scheme. More precisely, let $A_\nu \cong \mathbb{F}_\nu[\![\zeta]\!]$ be the completion of the local ring $\mathcal{O}_{C,\nu}$ at a closed point $\nu \in C$, let Q_ν be its fraction field, and let $\mathbb{P} = \mathbb{P}_\nu := \mathfrak{G} \times_C \operatorname{Spec} A_\nu$ and $P_\nu = \mathfrak{G} \times_C \operatorname{Spec} Q_\nu$. A *local* $\mathbb{P}_\nu$-*shtuka* over a scheme $S \in \mathcal{N}ilp_{A_\nu}$ is a pair $\underline{\mathcal{L}} = (\mathcal{L}_+, \hat{\tau})$ consisting of an $L^+\mathbb{P}_\nu$-torsor $\mathcal{L}_+$ on S and an isomorphism of the LP_ν-torsors $\hat{\tau} \colon \hat{\sigma}^* \mathcal{L} \xrightarrow{\sim} \mathcal{L}$. Here LP_ν (resp. $L^+\mathbb{P}_\nu$) denotes the group of loops (resp. positive loops) of $\mathbb{P}_\nu$ (see Section 2.1), $\mathcal{L}$ denotes the LP_ν-torsor associated with $\mathcal{L}_+$ and $\hat{\sigma}^* \mathcal{L}$ the pullback of $\mathcal{L}$ under the absolute $\mathbb{F}_\nu$-Frobenius endomorphism $\mathrm{Frob}_{(\#\mathbb{F}_\nu),S} \colon S \to S$. Moreover, $\mathcal{N}ilp_{A_\nu}$ denotes the category of A_ν-schemes on which the uniformizer ζ of A_ν is locally nilpotent. Local $\mathbb{P}_\nu$-shtukas can be viewed as function field analogs of p-divisible groups; see also [HV11, HV12].

As a preparation to [AH13] we show in this article that the Rapoport-Zink functor

$$\begin{array}{rcl} \underline{\mathcal{M}}_{\underline{\mathbb{L}}_0} \colon \mathcal{N}ilp_{k[\![\zeta]\!]} & \longrightarrow & \mathcal{S}ets \\ S & \longmapsto & \big\{ \text{Isomorphism classes of } (\underline{\mathcal{L}}, \bar{\delta}) \colon \text{ where } \underline{\mathcal{L}} \text{ is a local } \mathbb{P}_\nu\text{-shtuka} \\ & & \qquad \text{over } S \text{ and } \bar{\delta} \colon \underline{\mathcal{L}}_{\bar{S}} \to \underline{\mathbb{L}}_{0,\bar{S}} \text{ is a quasi-isogeny over } \bar{S} \big\}, \end{array}$$

for a fixed local $\mathbb{P}_\nu$-shtuka $\underline{\mathbb{L}}_0$ over a field k, and $\bar{S} = \mathrm{V}(\zeta) \subset S$, is representable by an ind-scheme, ind-quasi-projective over $\operatorname{Spf} k[\![\zeta]\!]$; see Theorem 4.4. To obtain a formal scheme locally formally of finite type, as in the analog for p-divisible groups, one has to bound the *Hodge polygon*, that is the relative position of $\hat{\sigma}^* \mathcal{L}_+$ and $\mathcal{L}_+$ under $\hat{\tau}$. We give an axiomatic treatment of bounds in Section 4.2 and prove the representability of the bounded Rapoport-Zink functor by a formal scheme locally formally of finite type over $\operatorname{Spf} k[\![\zeta]\!]$ in Theorem 4.13.

In addition, in Chapter 3 we discuss the relation between local $\mathbb{P}$-shtukas and Galois representations which is given by the associated Tate module. In Chapter 5 we consider the formal stack $\nabla_n \mathscr{H}^1(C, \mathfrak{G})^{\underline{\nu}}$, which is obtained by taking the formal completion of the stack $\nabla_n \mathscr{H}^1(C, \mathfrak{G})$ at a fixed n-tuple of pairwise different characteristic places $\underline{\nu} = (\nu_1, \ldots, \nu_n)$. This means we let $A_{\underline{\nu}}$ be the completion of the local ring $\mathcal{O}_{C^n,\underline{\nu}}$, and we consider global $\mathfrak{G}$-shtukas only over schemes S whose characteristic morphism $S \to C^n$ factors through $\mathcal{N}ilp_{A_{\underline{\nu}}}$. Recall that with an abelian variety over a scheme in $\mathcal{N}ilp_{\mathbb{Z}_p}$ one can associate its p-divisible group. In the analogous situation for global $\mathfrak{G}$-shtukas one can associate a tuple $(\widehat{\Gamma}_{\nu_i}(\underline{\mathcal{G}}))_i$ of local $\mathbb{P}_{\nu_i}$-shtukas $\widehat{\Gamma}_{\nu_i}(\underline{\mathcal{G}})$ with a global $\mathfrak{G}$-shtuka $\underline{\mathcal{G}}$ in $\nabla_n\mathscr{H}^1(C,\mathfrak{G})^{\underline{\nu}}(S)$. We construct this *global-local functor* in Section 5.2 by first generalizing the glueing lemma of Beauville and Laszlo [BL95] in Lemma 5.1. In analogy with a theorem of Serre and Tate relating the deformation theory of abelian varieties over schemes in $\mathcal{N}ilp_{\mathbb{Z}_p}$ and their associated p-divisible groups, we prove in Theorem 5.10 the equivalence between the infinitesimal deformations of a global $\mathfrak{G}$-shtuka and the infinitesimal deformations of its associated n-tuple of local $\mathbb{P}_{\nu_i}$-shtukas. Note that unlike abelian varieties, $\mathfrak{G}$-shtukas posses more than one characteristic and we must keep track of the deformations of the local $\mathbb{P}_{\nu_i}$-shtukas at each of these characteristic places ν_i. This theorem for abelian τ-sheaves (corresponding to the case $\mathfrak{G} = \mathrm{GL}_r$) and their associated z-divisible groups was first stated and proved by the second author in [Har05].

1.1 Notation and Conventions

Throughout this article we denote by

$\mathbb{F}_q$	a finite field with q elements and characteristic p,
C	a smooth projective geometrically irreducible curve over $\mathbb{F}_q$,
$Q := \mathbb{F}_q(C)$	the function field of C,
ν	a closed point of C, also called a *place* of C,
$\mathbb{F}_\nu$	the residue field at the place ν on C,
A_ν	the completion of the stalk $\mathcal{O}_{C,\nu}$ at ν,
$Q_\nu := \mathrm{Frac}(A_\nu)$	its fraction field,
$\mathbb{F}$	a finite field containing $\mathbb{F}_q$,
$\mathbb{D}_R := \operatorname{Spec} R[\![z]\!]$	the spectrum of the ring of formal power series in z with coefficients in an $\mathbb{F}$-algebra R,
$\hat{\mathbb{D}}_R := \operatorname{Spf} R[\![z]\!]$	the formal spectrum of $R[\![z]\!]$ with respect to the z-adic topology.

When $R = \mathbb{F}$ we drop the subscript R from the notation of $\mathbb{D}_R$ and $\hat{\mathbb{D}}_R$.

For a formal scheme $\widehat{S}$ we denote by $\mathcal{N}ilp_{\widehat{S}}$ the category of schemes over $\widehat{S}$ on which an ideal of definition of $\widehat{S}$ is locally nilpotent. We equip $\mathcal{N}ilp_{\widehat{S}}$ with the étale topology. We also denote by

$n \in \mathbb{N}_{>0}$	a positive integer,
$\underline{\nu} := (\nu_i)_{i=1\dots n}$	an n-tuple of closed points of C,
$A_{\underline{\nu}}$	the completion of the local ring $\mathcal{O}_{C^n,\underline{\nu}}$ of C^n at the closed point $\underline{\nu} = (\nu_i)$,
$\mathcal{N}ilp_{A_{\underline{\nu}}} := \mathcal{N}ilp_{\operatorname{Spf} A_{\underline{\nu}}}$	the category of schemes over C^n on which the ideal defining the closed point $\underline{\nu} \in C^n$ is locally nilpotent,
$\mathcal{N}ilp_{\mathbb{F}[\![\zeta]\!]} := \mathcal{N}ilp_{\hat{\mathbb{D}}}$	the category of $\mathbb{D}$-schemes S for which the image of z in $\mathcal{O}_S$ is locally nilpotent. We denote the image of z by ζ since we need to distinguish it from $z \in \mathcal{O}_{\mathbb{D}}$.
$\mathfrak{G}$	a parahoric Bruhat-Tits group scheme over C; see Definition 2.12,
$\mathbb{P}_\nu := \mathfrak{G} \times_C \operatorname{Spec} A_\nu$	the base change of $\mathfrak{G}$ to $\operatorname{Spec} A_\nu$,
$P_\nu := \mathfrak{G} \times_C \operatorname{Spec} Q_\nu$	the generic fiber of $\mathbb{P}_\nu$ over $\operatorname{Spec} Q_\nu$,
$\mathbb{P}$	a smooth affine group scheme of finite type over $\mathbb{D} = \operatorname{Spec} \mathbb{F}[\![z]\!]$,
$P := \mathbb{P} \times_{\mathbb{D}} \operatorname{Spec} \mathbb{F}(\!(z)\!)$	the generic fiber of $\mathbb{P}$ over $\operatorname{Spec} \mathbb{F}(\!(z)\!)$.

Let S be an $\mathbb{F}_q$-scheme. We denote by $\sigma_S\colon S\to S$ its $\mathbb{F}_q$-Frobenius endomorphism which acts as the identity on the points of S and as the q-power map on the structure sheaf. Likewise we let $\hat\sigma_S\colon S\to S$ be the $\mathbb{F}$-Frobenius endomorphism of an $\mathbb{F}$-scheme S. We set

$$C_S := C\times_{\operatorname{Spec}\mathbb{F}_q} S,\quad \text{and}$$
$$\sigma := \mathrm{id}_C\times\sigma_S.$$

Let H be a sheaf of groups (for the étale topology) on a scheme X. In this article a (*right*) *H-torsor* (also called an *H-bundle*) on X is a sheaf $\mathcal{G}$ for the étale topology on X together with a (right) action of the sheaf H such that $\mathcal{G}$ is isomorphic to H on a étale covering of X. Here H is viewed as an H-torsor by right multiplication.

2 Local ℙ-Shtukas and Global 𝔊-Shtukas

Global $\mathfrak{G}$-shtukas are function field analogs of abelian varieties. They were introduced by Drinfeld [Dri87] in the case where $\mathfrak{G}=\mathrm{GL}_r$ and used by him and by Lafforgue [Laf02] to establish the Langlands correspondence for GL_r over global function fields. We generalize Drinfeld's definition in Definition 2.14. The local p-adic properties of abelian varieties are largely captured by their associated p-divisible groups. In the theory of global $\mathfrak{G}$-shtukas the latter correspond to local shtukas.

2.1 Loop Groups and Local ℙ-Shtukas

Since we want to develop the theory of local $\mathbb{P}$-shtukas partly independently of global $\mathfrak{G}$-shtukas we let $\mathbb{F}$ be a finite field and $\mathbb{F}[\![z]\!]$ be the power series ring over $\mathbb{F}$ in the variable z. We let $\mathbb{P}$ be a smooth affine group scheme over $\mathbb{D}:=\operatorname{Spec}\mathbb{F}[\![z]\!]$, and we let $P:=\mathbb{P}\times_{\mathbb{D}}\dot{\mathbb{D}}$ be the generic fiber of $\mathbb{P}$ over $\dot{\mathbb{D}}:=\operatorname{Spec}\mathbb{F}(\!(z)\!)$.

Definition 2.1. The *group of positive loops associated with* $\mathbb{P}$ is the infinite dimensional affine group scheme $L^+\mathbb{P}$ over $\mathbb{F}$ whose R-valued points for an $\mathbb{F}$-algebra R are

$$L^+\mathbb{P}(R):=\mathbb{P}(R[\![z]\!]):=\mathbb{P}(\mathbb{D}_R):=\operatorname{Hom}_{\mathbb{D}}(\mathbb{D}_R,\mathbb{P})\,.$$

The *group of loops associated with* P is the *fpqc*-sheaf of groups LP over $\mathbb{F}$ whose R-valued points for an $\mathbb{F}$-algebra R are

$$LP(R):=P(R(\!(z)\!)):=P(\dot{\mathbb{D}}_R):=\operatorname{Hom}_{\dot{\mathbb{D}}}(\dot{\mathbb{D}}_R,P)\,,$$

where we write $R(\!(z)\!):=R[\![z]\!][\frac{1}{z}]$ and $\dot{\mathbb{D}}_R:=\operatorname{Spec}R(\!(z)\!)$. Let $\mathscr{H}^1(\operatorname{Spec}\mathbb{F},L^+\mathbb{P}):=[\operatorname{Spec}\mathbb{F}/L^+\mathbb{P}]$ (resp. $\mathscr{H}^1(\operatorname{Spec}\mathbb{F},LP):=[\operatorname{Spec}\mathbb{F}/LP]$) denote the classifying space of $L^+\mathbb{P}$-torsors (resp. LP-torsors). It is a stack fibered in groupoids over the category of $\mathbb{F}$-schemes S whose category $\mathscr{H}^1(\operatorname{Spec}\mathbb{F},L^+\mathbb{P})(S)$ consists of all $L^+\mathbb{P}$-torsors (resp. LP-torsors) on S. The inclusion of sheaves $L^+\mathbb{P}\subset LP$ gives rise to the natural 1-morphism

$$L\colon \mathscr{H}^1(\operatorname{Spec}\mathbb{F},L^+\mathbb{P})\longrightarrow\mathscr{H}^1(\operatorname{Spec}\mathbb{F},LP),\ \mathcal{L}_+\mapsto\mathcal{L}\,. \tag{2.1}$$

Definition 2.2. Let $\hat{\mathbb{P}}$ be the formal group scheme over $\hat{\mathbb{D}}:=\operatorname{Spf}\mathbb{F}[\![z]\!]$, obtained by the formal completion of $\mathbb{P}$ along $V(z)$. A *formal $\hat{\mathbb{P}}$-torsor* over an $\mathbb{F}$-scheme S is a z-adic formal scheme $\hat{\mathcal{P}}$ over $\hat{\mathbb{D}}_S:=\hat{\mathbb{D}}\,\widehat{\times}_{\mathbb{F}}\,S$ together with an action $\hat{\mathbb{P}}\,\widehat{\times}_{\hat{\mathbb{D}}}\,\hat{\mathcal{P}}\to\hat{\mathcal{P}}$ of $\hat{\mathbb{P}}$ on $\hat{\mathcal{P}}$ such that there is a covering $\hat{\mathbb{D}}_{S'}\to\hat{\mathbb{D}}_S$ where $S'\to S$ is an *fpqc*-covering and a $\hat{\mathbb{P}}$-equivariant isomorphism

$$\hat{\mathcal{P}}\,\widehat{\times}_{\hat{\mathbb{D}}_S}\,\hat{\mathbb{D}}_{S'}\xrightarrow{\sim}\hat{\mathbb{P}}\,\widehat{\times}_{\hat{\mathbb{D}}}\,\hat{\mathbb{D}}_{S'}.$$

Here $\hat{\mathbb{P}}$ acts on itself by right multiplication. Let $\mathscr{H}^1(\hat{\mathbb{D}},\hat{\mathbb{P}})$ be the category fibered in groupoids that assigns to each $\mathbb{F}$-scheme S the groupoid consisting of all formal $\hat{\mathbb{P}}$-torsors over $\hat{\mathbb{D}}_S$.

Remark 2.3. For any $\widehat{\mathcal{P}}$ in $\mathscr{H}^1(\hat{\mathbb{D}},\hat{\mathbb{P}})(\operatorname{Spec}R)$ one can find an étale covering $R\to R'$ such that $\widehat{\mathcal{P}}\,\hat{\times}_{\hat{\mathbb{D}}_R}\hat{\mathbb{D}}_{R'}$ is isomorphic to $\hat{\mathbb{P}}_{R'}$ in $\mathscr{H}^1(\hat{\mathbb{D}},\hat{\mathbb{P}})(R')$. Indeed, since $\widehat{\mathcal{P}}\to\hat{\mathbb{D}}_R$ is smooth, the restriction $\widehat{\mathcal{P}}_0$ of $\widehat{\mathcal{P}}$ to $V(z)\subseteq\hat{\mathbb{D}}_R$ is likewise smooth over R. Therefore $\widehat{\mathcal{P}}_0$ has a section over an étale covering $R\to R'$. Then by smoothness this section extends over $\hat{\mathbb{D}}_R$.

In [HV11, Proposition 2.2.(a)] Viehmann and the second author proved that for a split reductive group G, there is a bijection of (pointed) sets between the Čech cohomology $\check{\mathrm{H}}^1(S_{fpqc},L^+G)$ and the set of isomorphism classes of z-adic formal schemes over $\hat{\mathbb{D}}_S$. By the same arguments one can even see that there is a canonical equivalence between the corresponding categories.

Proposition 2.4. *There is a natural isomorphism*

$$\mathscr{H}^1(\hat{\mathbb{D}},\hat{\mathbb{P}})\xrightarrow{\sim}\mathscr{H}^1(\operatorname{Spec}\mathbb{F},L^+\mathbb{P})$$

of groupoids. In particular all $L^+\mathbb{P}$-torsors for the fpqc-topology on S are already trivial étale locally on S.

Proof. With a given element $\hat{\mathcal{P}}$ of $\mathscr{H}^1(\hat{\mathbb{D}},\hat{\mathbb{P}})(S)$ one can associate the following sheaf

$$\begin{aligned}\underline{\mathcal{K}}\colon\ S_{fpqc}&\longrightarrow\mathcal{S}ets\\ T&\longmapsto\operatorname{Hom}_{\hat{\mathbb{D}}_S}(\hat{\mathbb{D}}_T,\hat{\mathcal{P}}),\end{aligned}$$

where S_{fpqc} denotes the big *fpqc*-site on S. This sheaf is a torsor under the action of $L^+\mathbb{P}(T)=\operatorname{Hom}_{\hat{\mathbb{D}}}(\hat{\mathbb{D}}_T,\hat{\mathbb{P}})$.

Conversely let $\mathcal{K}$ be an $L^+\mathbb{P}$-torsor. Let $S'\to S$ be a covering that trivializes $\mathcal{K}$ and fix a trivialization $\mathcal{K}_{S'}\xrightarrow{\sim}(L^+\mathbb{P})_{S'}$. This gives a 1-cocycle $g\in L^+\mathbb{P}(S'')$, where $S''=S'\times_S S'$. Now $\bar{g}=g(mod\ z^n)$ can be viewed as a descent data on $\hat{\mathbb{P}}\,\hat{\times}_{\mathbb{D}}\,\mathbb{D}_{n,S'}=\mathbb{P}\times_{\mathbb{D}}\mathbb{D}_{n,S'}$ where $\mathbb{D}_{n,S'}:=\operatorname{Spec}\mathbb{F}[\![z]\!]/(z^n)\times_{\mathbb{F}}S'$. Since $\mathbb{D}_{n,S'}\to\mathbb{D}_{n,S}$ is an *fpqc*-covering and $\mathbb{P}$ is affine, the descent data is effective by [BLR90, §6.1, Theorem 6] and gives an affine finitely presented smooth scheme $\hat{\mathcal{G}}_n$ over $\mathbb{D}_{n,S}$ by [EGA, IV_2, Proposition 2.7.1 and IV_4, Corollaire 17.7.3]. These schemes form an inductive system $\{\hat{\mathcal{G}}_n\}_{n\in\mathbb{N}}$. Now set $\hat{\mathcal{G}}:=\varinjlim\hat{\mathcal{G}}_n$, the existence of this limit (in the category of z-adic formal schemes over $\hat{\mathbb{D}}_S$) follows from [EGA, $\mathrm{I}_{\mathrm{new}}$, Corollary 10.6.4]. This shows that the functor is essentially surjective. By the above construction we see that the functor is also fully faithful.

The last statement now follows from Remark 2.3. □

Definition 2.5. Assume that we have two morphisms $f,g\colon X\to Y$ of schemes or stacks. We denote by $\operatorname{equi}(f,g\colon X\rightrightarrows Y)$ the pull back of the diagonal under the morphism $(f,g)\colon X\to Y\times_{\mathbb{Z}}Y$, that is, we let $\operatorname{equi}(f,g\colon X\rightrightarrows Y):=X\times_{(f,g),Y\times Y,\Delta}Y$ where $\Delta=\Delta_{Y/\mathbb{Z}}\colon Y\to Y\times_{\mathbb{Z}}Y$ is the diagonal morphism.

We define the space of local $\mathbb{P}$-shtukas as follows.

Definition 2.6. Let $\mathcal{X}$ be the fiber product

$$\mathscr{H}^1(\operatorname{Spec}\mathbb{F},L^+\mathbb{P})\times_{\mathscr{H}^1(\operatorname{Spec}\mathbb{F},LP)}\mathscr{H}^1(\operatorname{Spec}\mathbb{F},L^+\mathbb{P})$$

of groupoids. Let pr_i denote the projection onto the i-th factor. We define the groupoid of *local $\mathbb{P}$-shtukas* $Sht_{\mathbb{P}}^{\mathbb{D}}$ to be

$$Sht_{\mathbb{P}}^{\mathbb{D}}\ :=\ \operatorname{equi}\left(\hat{\sigma}\circ pr_1,pr_2\colon\mathcal{X}\rightrightarrows\mathscr{H}^1(\operatorname{Spec}\mathbb{F},L^+\mathbb{P})\right)\hat{\times}_{\operatorname{Spec}\mathbb{F}}\operatorname{Spf}\mathbb{F}[\![\zeta]\!].$$

(see Definition 2.5) where $\hat{\sigma} := \hat{\sigma}_{\mathscr{H}^1(\operatorname{Spec}\mathbb{F}, L^+\mathbb{P})}$ is the $\mathbb{F}$-Frobenius of $\mathscr{H}^1(\operatorname{Spec}\mathbb{F}, L^+\mathbb{P})$. The category $Sht_{\mathbb{P}}^{\mathbb{D}}$ is fibered in groupoids over the category $\mathcal{N}ilp_{\mathbb{F}[\![\zeta]\!]}$ of $\mathbb{F}[\![\zeta]\!]$-schemes on which ζ is locally nilpotent. We call an object of the category $Sht_{\mathbb{P}}^{\mathbb{D}}(S)$ a *local $\mathbb{P}$-shtuka over S.*

More explicitly a local $\mathbb{P}$-shtuka over $S \in \mathcal{N}ilp_{\mathbb{F}[\![\zeta]\!]}$ is a pair $\underline{\mathcal{L}} = (\mathcal{L}_+, \hat{\tau})$ consisting of an $L^+\mathbb{P}$-torsor $\mathcal{L}_+$ on S and an isomorphism of the associated loop group torsors $\hat{\tau}\colon \hat{\sigma}^*\mathcal{L} \to \mathcal{L}$ from (2.1).

Definition 2.7. A local $\mathbb{P}$-shtuka $(\mathcal{L}_+, \hat{\tau})$ is called *étale* if $\hat{\tau}$ comes from an isomorphism of $L^+\mathbb{P}$-torsors $\hat{\sigma}^*\mathcal{L}_+ \xrightarrow{\sim} \mathcal{L}_+$.

Lemma 2.8. *Let k be an algebraically closed field extension of $\mathbb{F}$. Then for any $b \in L^+\mathbb{P}(k)$ there exists some $c \in L^+\mathbb{P}(k)$ such that $b\,\hat{\sigma}^*(c) = c$.*

Proof. Let $\hat{\mathbb{P}}$ be as in Definition 2.2. Then $L^+\mathbb{P}(k) = \hat{\mathbb{P}}(k[\![z]\!])$. We view $\hat{\mathbb{P}}$ as the inductive limit $\varinjlim \mathbb{P}_n$, where $\mathbb{P}_n = \mathbb{P} \times_{\mathbb{D}} \mathbb{D}_n$ with $\mathbb{D}_n := \operatorname{Spec}\mathbb{F}[\![z]\!]/(z^n)$. Let $\widetilde{G}_n$ denote the linear algebraic group over $\mathbb{F}$ given by the Weil restriction $\operatorname{Res}_{\mathbb{D}_n/\operatorname{Spec}\mathbb{F}}(\mathbb{P}_n)$. The reduction of b mod z^n gives an element $b_n \in \widetilde{G}_n(k)$. By Lang's theorem [Lan56, Corollary on p. 557] there exist a $c_n \in \widetilde{G}_n(k)$ such that $b_n\hat{\sigma}^*(c_n) = c_n$. Here $\hat{\sigma}$ is the $\mathbb{F}$-Frobenius on $\widetilde{G}_n$ which coincides with the Frobenius $\hat{\sigma}$ induced from $\mathbb{P}$. Now consider the reduction map $\alpha_n\colon \widetilde{G}_{n+1}(k) \to \widetilde{G}_n(k)$ and the element $\bar{d}_n := \alpha_n(c_{n+1})^{-1}c_n$ which satisfies $\hat{\sigma}^*(\bar{d}_n) = \bar{d}_n$ and hence lies in $\widetilde{G}_n(\mathbb{F}) = \mathbb{P}(\mathbb{D}_{n+1})$. Since $\mathbb{P}$ is smooth $\bar{d}_n$ lifts to an element $d_n \in \mathbb{P}(\mathbb{D}_{n+1}) = \widetilde{G}_{n+1}(\mathbb{F})$. Replacing c_{n+1} by $c_{n+1}d_n$ we may assume that $\alpha_n(c_{n+1}) = c_n$ and then take $c := \varprojlim c_n$. □

Corollary 2.9. *Every étale local $\mathbb{P}$-shtuka over an algebraically closed field k is isomorphic to $\big((L^+\mathbb{P})_k, 1\cdot\hat{\sigma}^*\big)$.*

Proof. Let $\underline{\mathcal{L}} = (\mathcal{L}_+, \hat{\tau})$ be an étale local $\mathbb{P}$-shtuka over k. By Proposition 2.4 there is a trivialization of the $L^+\mathbb{P}$-torsor $\mathcal{L}_+$ giving rise to an isomorphism $\underline{\mathcal{L}} \cong \big((L^+\mathbb{P})_k, b\cdot\hat{\sigma}^*\big)$ for some $b \in L^+\mathbb{P}(k)$. By Lemma 2.8 there is an element $c \in L^+\mathbb{P}(k)$ with $b\,\hat{\sigma}^*(c) = c$ and multiplication with c is an isomorphism $\big((L^+\mathbb{P})_k, 1\cdot\hat{\sigma}^*\big) \xrightarrow{\sim} \big((L^+\mathbb{P})_k, b\cdot\hat{\sigma}^*\big)$. □

Local $\mathbb{P}$-shtukas can be viewed as function field analogs of p-divisible groups. This inspires the following notions of quasi-isogenies.

Definition 2.10. A *quasi-isogeny* $f\colon \underline{\mathcal{L}} \to \underline{\mathcal{L}}'$ between two local $\mathbb{P}$-shtukas $\underline{\mathcal{L}} := (\mathcal{L}_+, \hat{\tau})$ and $\underline{\mathcal{L}}' := (\mathcal{L}'_+, \hat{\tau}')$ over S is an isomorphism of the associated LP-torsors $f\colon \mathcal{L} \to \mathcal{L}'$ satisfying $f \circ \hat{\tau} = \hat{\tau}' \circ \hat{\sigma}^* f$. We denote by $\operatorname{QIsog}_S(\underline{\mathcal{L}}, \underline{\mathcal{L}}')$ the set of quasi-isogenies between $\underline{\mathcal{L}}$ and $\underline{\mathcal{L}}'$ over S, and we write $\operatorname{QIsog}_S(\underline{\mathcal{L}}) := \operatorname{QIsog}_S(\underline{\mathcal{L}}, \underline{\mathcal{L}})$ for the quasi-isogeny group of $\underline{\mathcal{L}}$.

As in the theory of p-divisible groups, also our quasi-isogenies are rigid.

Proposition 2.11 (Rigidity of quasi-isogenies for local $\mathbb{P}$-shtukas)**.** *Let S be a scheme in $\mathcal{N}ilp_{\mathbb{F}[\![\zeta]\!]}$ and let $j\colon \bar{S} \to S$ be a closed immersion defined by a sheaf of ideals $\mathcal{I}$ which is locally nilpotent. Let $\underline{\mathcal{L}}$ and $\underline{\mathcal{L}}'$ be two local $\mathbb{P}$-shtukas over S. Then*

$$\operatorname{QIsog}_S(\underline{\mathcal{L}}, \underline{\mathcal{L}}') \longrightarrow \operatorname{QIsog}_{\bar{S}}(j^*\underline{\mathcal{L}}, j^*\underline{\mathcal{L}}'), \quad f \mapsto j^*f$$

is a bijection of sets.

Proof. This was proved in [HV11, Proposition 3.9] when $\mathbb{P} = G \times_{\mathbb{F}} \mathbb{D}$ for a constant split reductive group G over $\mathbb{F}$. The proof carries over literally. Compare also Proposition 5.9. □

2.2 Global $\mathfrak{G}$-Shtukas

Let $\mathbb{F}_q$ be a finite field with q elements, let C be a smooth projective geometrically irreducible curve over $\mathbb{F}_q$, and let $\mathfrak{G}$ be a parahoric group scheme over C as in the following

Definition 2.12. A smooth affine group scheme $\mathfrak{G}$ over C is called a *parahoric* (*Bruhat-Tits*) *group scheme* if

(a) all geometric fibers of $\mathfrak{G}$ are connected and the generic fiber of $\mathfrak{G}$ is reductive over $\mathbb{F}_q(C)$,

(b) for any ramification point ν of $\mathfrak{G}$ (i.e. those points ν of C, for which the fiber above ν is not reductive) the group scheme $\mathbb{P}_\nu := \mathfrak{G} \times_C \operatorname{Spec} A_\nu$ is a parahoric group scheme over A_ν, as defined by Bruhat and Tits [BT72, Définition 5.2.6]; see also [HR03].

The relation to Section 2.1 is as follows. We are mainly interested in the case where $\mathbb{F}[\![z]\!] \cong A_\nu$ and $\mathbb{P} = \mathbb{P}_\nu$.

Proposition 2.13. *Let $\mathbb{P}$ be a parahoric group scheme over $\mathbb{D} = \operatorname{Spec} \mathbb{F}[\![z]\!]$ in the sense of [BT72, Définition 5.2.6] and [HR03].*

(a) There is a faithful representation $\mathbb{P} \hookrightarrow \mathrm{SL}_{r,\mathbb{D}}$ over $\mathbb{D}$ with quasi-affine quotient $\mathrm{SL}_{r,\mathbb{D}} / \mathbb{P}$,

(b) If all fibers of $\mathbb{P}$ over $\mathbb{D}$ are reductive then there exists a closed immersion of group schemes $\mathbb{P} \hookrightarrow \mathrm{SL}_{r,\mathbb{D}}$ with affine quotient $\mathrm{SL}_{r,\mathbb{D}} / \mathbb{P}$.

Proof. Pappas and Rapoport [PR08, Proposition 1.3] show that there is a closed immersion $\mathbb{P} \hookrightarrow \mathrm{GL}_{r,\mathbb{D}} \times_{\mathbb{D}} \mathbb{G}_{m,\mathbb{D}} =: G$ with quasi-affine quotient $G/\mathbb{P}$. We choose a closed immersion $G \hookrightarrow \mathrm{SL}_{r',\mathbb{D}}$ and use that the property of being (quasi-)affine is *fpqc*-local on the base by [EGA, Proposition 2.7.1]. Then $\mathrm{SL}_{r',\mathbb{D}} / G$ is affine by Haboush's theorem [Hab78, Theorem 3.3]. Also the morphism $\mathrm{SL}_{r',\mathbb{D}} / \mathbb{P} \to \mathrm{SL}_{r',\mathbb{D}} / G$ is quasi-affine because after base change to $\mathrm{SL}_{r',\mathbb{D}}$ it equals the projection $(G/\mathbb{P}) \times_{\mathbb{D}} \mathrm{SL}_{r',\mathbb{D}} \to \mathrm{SL}_{r',\mathbb{D}}$. This proves (a).

By [EGA, II, Proposition 5.1.2] the question whether a quasi-affine morphism is affine can be tested fiber-wise. So (b) follows from Haboush's theorem [Hab78, Theorem 3.3] again. □

Definition 2.14. A global $\mathfrak{G}$-shtuka $\underline{\mathcal{G}}$ over an $\mathbb{F}_q$-scheme S is a tuple $(\mathcal{G}, s_1, \ldots, s_n, \tau)$ where

$\mathcal{G}$ is a $\mathfrak{G}$-torsor $\mathcal{G}$ over C_S,

$s_1, \ldots, s_n \in C(S)$ are $\mathbb{F}_q$-morphisms called the *characteristic sections of* $\underline{\mathcal{G}}$, and

$\tau \colon \sigma^*\mathcal{G}_{|C_S \smallsetminus \Gamma_{s_1} \cup \ldots \cup \Gamma_{s_n}} \xrightarrow{\sim} \mathcal{G}_{|C_S \smallsetminus \Gamma_{s_1} \cup \ldots \cup \Gamma_{s_n}}$ is an isomorphism of $\mathfrak{G}$-torsors. Here $\Gamma_{s_i} \subset C_S$ denotes the graph of the morphisms s_i.

Here $\sigma = \mathrm{id}_C \times \sigma_S$ for the $\mathbb{F}_q$-Frobenius endomorphism $\sigma_S \colon S \to S$ which acts as the identity on the points of S and as the q-power map on the structure sheaf. We denote the *moduli stack* of global $\mathfrak{G}$-shtukas by $\nabla_n \mathscr{H}^1(C, \mathfrak{G})$. It is a stack fibered in groupoids over the category of $\mathbb{F}_q$-schemes. Sometimes we will fix the sections $(s_1, \ldots, s_n) \in C^n(S)$ and simply call $\underline{\mathcal{G}} = (\mathcal{G}, \tau)$ a global $\mathfrak{G}$-shtuka over S.

Although the definition of global $\mathfrak{G}$-shtuka makes sense also if $\mathfrak{G}$ is just a smooth affine group scheme over C, we require that $\mathfrak{G}$ is parahoric in [AH13, Theorem 3.14] where we prove the following

Theorem 2.15. *The stack $\nabla_n \mathscr{H}_D^1(C, \mathfrak{G})$ is an ind-algebraic stack (in the sense of [AH13, Definition 3.13]) over C^n which is ind-separated and locally of ind-finite type.*

There is also a notion of quasi-isogenies for global $\mathfrak{G}$-shtukas.

Definition 2.16. Consider a scheme S together with characteristic morphisms $s_i\colon S \to C$ for $i = 1, \dots, n$ and let $\underline{\mathcal{G}} = (\mathcal{G}, \tau)$ and $\underline{\mathcal{G}}' = (\mathcal{G}', \tau')$ be two global $\mathfrak{G}$-shtukas over S with the same characteristics s_i. A *quasi-isogeny* from $\underline{\mathcal{G}}$ to $\underline{\mathcal{G}}'$ is an isomorphism $f\colon \mathcal{G}|_{C_S \smallsetminus D_S} \xrightarrow{\sim} \mathcal{G}'|_{C_S \smallsetminus D_S}$ satisfying $\tau'\sigma^*(f) = f\tau$, where D is some effective divisor on C. We denote the *group of quasi-isogenies* of $\underline{\mathcal{G}}$ to itself by $\operatorname{QIsog}_S(\underline{\mathcal{G}})$.

Like for abelian varieties, rigidity of global $\mathfrak{G}$-shtukas only holds in fixed finite characteristics; see Proposition 5.9.

3 Tate Modules for Local ℙ-Shtukas

In this chapter we assume that $\mathbb{P}$ is a smooth affine group scheme over $\mathbb{D}$. For a scheme $S \in \mathcal{N}ilp_{\mathbb{F}[\![\zeta]\!]}$ let $\mathcal{O}_S[\![z]\!]$ be the sheaf of $\mathcal{O}_S$-algebras on S for the *fpqc*-topology whose ring of sections on an S-scheme Y is the ring of power series $\mathcal{O}_S[\![z]\!](Y) := \Gamma(Y, \mathcal{O}_Y)[\![z]\!]$. Let $\mathcal{O}_S(\!(z)\!)$ be the *fpqc*-sheaf of $\mathcal{O}_S$-algebras on S associated with the presheaf $Y \mapsto \Gamma(Y, \mathcal{O}_Y)[\![z]\!][\frac{1}{z}]$. A sheaf M of $\mathcal{O}_S[\![z]\!]$-modules on S which is free *fpqc*-locally on S is already free Zariski-locally on S by [HV11, Proposition 2.3]. We call those modules *locally free sheaves of* $\mathcal{O}_S[\![z]\!]$*-modules.* We denote by $\hat{\sigma}^*$ the endomorphism of $\mathcal{O}_S[\![z]\!]$ and $\mathcal{O}_S(\!(z)\!)$ that acts as the identity on the variable z, and is the $\mathbb{F}$-Frobenius $b \mapsto (b)^{\#\mathbb{F}}$ on local sections $b \in \mathcal{O}_S$. For a sheaf M of $\mathcal{O}_S[\![z]\!]$-modules on S we set $\hat{\sigma}^* M := M \otimes_{\mathcal{O}_S[\![z]\!], \hat{\sigma}^*} \mathcal{O}_S[\![z]\!]$.

Definition 3.1. (a) A *local shtuka over* S is a pair $(M, \hat{\tau})$ where M is a locally free sheaf of $\mathcal{O}_S[\![z]\!]$-modules of finite rank on S and $\hat{\tau}\colon \hat{\sigma}^* M \otimes_{\mathcal{O}_S[\![z]\!]} \mathcal{O}_S(\!(z)\!) \xrightarrow{\sim} M \otimes_{\mathcal{O}_S[\![z]\!]} \mathcal{O}_S(\!(z)\!)$ is an isomorphism.

(b) A local shtuka $\underline{M} := (M, \hat{\tau})$ is called *étale* if $\hat{\tau}$ comes from an isomorphism of $\mathcal{O}_S[\![z]\!]$-modules $\hat{\sigma}^* M \xrightarrow{\sim} M$.

(c) A *morphism* $f\colon (M, \hat{\tau}) \to (M', \hat{\tau}')$ of local shtukas over S is a morphism $f\colon M \to M'$ of $\mathcal{O}_S[\![z]\!]$-modules which satisfies $\hat{\tau}' \circ \hat{\sigma}^* f = f \circ \hat{\tau}$. We do not require that f is an isomorphism. We denote by $Sht_{\mathbb{D}}(S)$ the category of local shtukas over S.

Remark 3.2. There is an equivalence of categories between $\mathscr{H}^1(\operatorname{Spec} \mathbb{F}, L^+ \mathrm{GL}_r)(S)$ and the category of locally free sheaves of $\mathcal{O}_S[\![z]\!]$-modules of rank r; see [HV11, §4]. It induces an equivalence between the category of local GL_r-shtukas over S and the category consisting of local shtukas over S of rank r with isomorphisms as the only morphisms; see [HV11, Lemma 4.2].

Definition 3.3. A *quasi-isogeny* between two local shtukas $(M, \hat{\tau}) \to (M', \hat{\tau}')$ is an isomorphism of $\mathcal{O}_S(\!(z)\!)$-modules

$$f\colon M \otimes_{\mathcal{O}_S[\![z]\!]} \mathcal{O}_S(\!(z)\!) \xrightarrow{\sim} M' \otimes_{\mathcal{O}_S[\![z]\!]} \mathcal{O}_S(\!(z)\!)$$

with $\hat{\tau}' \circ \hat{\sigma}^* f = f \circ \hat{\tau}$.

In analogy with p-divisible groups and abelian varieties, one can also assign a Galois representation to a given local shtuka as follows. Assume that S is connected. Let $\bar{s}$ be a geometric point of S and let $\pi_1(S, \bar{s})$ denote the algebraic fundamental group of S at $\bar{s}$. We define the (*dual*) *Tate functor* from the category of local shtukas $Sht_{\mathbb{D}}(S)$ over S to the category $\mathfrak{FMod}_{\mathbb{F}[\![z]\!][\pi_1(S,\bar{s})]}$ of $\mathbb{F}[\![z]\!][\pi_1(S, \bar{s})]$-modules which are finite free over $\mathbb{F}[\![z]\!]$ as follows

$$\begin{aligned} \check{T}_{-} : Sht_{\mathbb{D}}(S) &\longrightarrow \mathfrak{FMod}_{\mathbb{F}[\![z]\!][\pi_1(S,\bar{s})]}, \\ \underline{M} := (M, \hat{\tau}) &\longmapsto \check{T}_{\underline{M}} := (M \otimes_{\mathcal{O}_S[\![z]\!]} \kappa(\bar{s})[\![z]\!])^{\hat{\tau}}. \end{aligned}$$

Here the superscript $\hat\tau$ denotes $\hat\tau$-invariants. Sometimes also the notation $\mathrm{H}^1_{\acute{e}t}(\underline{M}, \mathbb{F}[\![z]\!]) := \check{T}_{\underline{M}}$ is used. Inverting z we also consider the *rational (dual) Tate functor*

$$\begin{aligned} \check V_- \colon Sht_{\mathbb D}(S) &\longrightarrow \mathfrak{FMod}_{\mathbb F(\!(z)\!)[\pi_1(S,\bar s)]},\\ \underline M := (M,\hat\tau) &\longmapsto \check V_{\underline M} := \big(M\otimes_{\mathcal O_S[\![z]\!]}\kappa(\bar s)[\![z]\!]\big)^{\hat\tau}\otimes_{\mathbb F[\![z]\!]}\mathbb F(\!(z)\!).\end{aligned}$$

where $\mathfrak{FMod}_{\mathbb F(\!(z)\!)[\pi_1(S,\bar s)]}$ denotes the category of $\mathbb F(\!(z)\!)[\pi_1(S,\bar s)]$-modules which are finite over $\mathbb F(\!(z)\!)$. The functor $\check V_-$ transforms quasi-isogenies into isomorphisms.

Proposition 3.4. *Let $S \in \mathcal{N}ilp_{\mathbb F[\![\zeta]\!]}$ be connected. Then the functor $\check T_-$ is an equivalence between the category of* étale *local shtukas over S and the category $\mathfrak{FMod}_{\mathbb F[\![z]\!][\pi_1(S,\bar s)]}$. The functor $\check V_-$ is an equivalence between the category of* étale *local shtukas over S with quasi-isogenies as morphisms and the category $\mathfrak{FMod}_{\mathbb F(\!(z)\!)[\pi_1(S,\bar s)]}$ with isomorphisms as the only morphisms.*

Proof. The statement for $\check T_-$ follows from the argument in [Har11, Proposition 1.3.7]. It is analogous to [Kat73, Proposition 4.1.1] and can be thought of as a positive characteristic analog of the Riemann-Hilbert correspondence. The quasi-inverse functor is explicitly constructed as follows. Consider an $\mathbb F[\![z]\!][\pi_1(S,\bar s)]$-module of rank r and the corresponding representation $\pi\colon \pi_1(S,\bar s)\to \mathrm{GL}_r(\mathbb F[\![z]\!])$. For each $m\in\mathbb N$ let $S_m\to S$ be the finite Galois covering corresponding to the kernel of $\pi_1(S,\bar s)\xrightarrow{\pi} \mathrm{GL}_r(\mathbb F[\![z]\!])\xrightarrow{\bmod z^m}\mathrm{GL}_r\big(\mathbb F[\![z]\!]/(z^m)\big)$. Let $\widetilde M_m$ be the free module of rank r over $\mathcal O_{S_m}[\![z]\!]/(z^m) = \mathcal O_{S_m}\otimes_{\mathbb F}\mathbb F[\![z]\!]/(z^m)$ and equip it with the Frobenius $\hat\tau := \hat\sigma\otimes\mathrm{id}$ and the action of $\gamma\in\mathrm{Gal}(S_m/S)$ by $\gamma(b\otimes f) = \gamma^*(b)\otimes\pi(\gamma^{-1})(f)$ for $b\in\mathcal O_{S_m}$ and $f\in\mathbb F[\![z]\!]/(z^m)^{\oplus r}$. Then $\widetilde M_m$ descends to a locally free $\mathcal O_S[\![z]\!]/(z^m)$-module M_m of rank r and $\hat\tau$ descends to an isomorphism $\hat\tau\colon\hat\sigma^*M_m\xrightarrow{\sim}M_m$. This makes $M := \varprojlim M_m$ into an étale local shtuka over S and yields the quasi-inverse to $\check T_-$.

That $\check V_-$ is essentially surjective follows from the fact that $\pi_1(S,\bar s)$ is compact which implies that every $\mathbb F(\!(z)\!)[\pi_1(S,\bar s)]$-module arises from an $\mathbb F[\![z]\!][\pi_1(S,\bar s)]$-module by inverting z. To see that $\check V_-$ is fully faithful consider two local shtukas $\underline M$ and $\underline M'$ over S and an isomorphism $f\colon \check V_{\underline M}\xrightarrow{\sim}\check V_{\underline M'}$. There are powers z^N and $z^{N'}$ of z such that z^Nf and $z^{N'}f^{-1}$ come from morphisms $\check T_{\underline M}\to\check T_{\underline M'}$ respectively $\check T_{\underline M'}\to\check T_{\underline M}$. Under the equivalence $\check T_-$ these in turn come from morphisms $g\colon\underline M\to\underline M'$ and $g'\colon\underline M'\to\underline M$. Then $gg' = z^{N+N'}$ and this implies that g and g' are quasi-isogenies. Clearly $\check V_{z^{-N}g} = f$ and $\check V_{z^{-N'}g'} = f^{-1}$. This proves that $\check V_-$ is an equivalence of categories. □

Let $Vect_{\mathbb D}$ be the groupoid over $\mathcal{N}ilp_{\mathbb F[\![\zeta]\!]}$ whose S-valued points is the category of locally free sheaves of $\mathcal O_S[\![z]\!]$-modules with isomorphisms as the only morphisms. Let $\mathrm{Rep}_{\mathbb F[\![z]\!]}\,\mathbb P$ be the category of representations $\rho\colon\mathbb P\to\mathrm{GL}(V)$ of $\mathbb P$ in finite free $\mathbb F[\![z]\!]$-modules V, that is, ρ is a morphism of algebraic groups over $\mathbb F[\![z]\!]$. Any such representation ρ gives a functor

$$\rho_*\colon \mathscr H^1(\operatorname{Spec}\mathbb F, L^+\mathbb P)\to Vect_{\mathbb D}$$

which sends an $L^+\mathbb P$-torsor $\mathcal L_+$ to the sheaf of $\mathcal O_S[\![z]\!]$-modules associated with the following presheaf

$$Y \longmapsto \Big(\mathcal L_+(Y)\times\big(V\otimes_{\mathbb F[\![z]\!]}\mathcal O_S[\![z]\!](Y)\big)\Big)/L^+\mathbb P(Y)\,. \tag{3.1}$$

The functor $\rho_*\colon\mathscr H^1(\operatorname{Spec}\mathbb F, L^+\mathbb P)\to Vect_{\mathbb D}$ induces a functor from the category of local $\mathbb P$-shtukas to the category of local shtukas which we likewise denote ρ_*. This functor is also compatible with quasi-isogenies.

Definition 3.5. Let $Funct^{\otimes}(\mathrm{Rep}_{\mathbb{F}[\![z]\!]}\,\mathbb{P}, \mathfrak{FMod}_{\mathbb{F}[\![z]\!][\pi_1(S,\bar{s})]})$, resp. $Funct^{\otimes}(\mathrm{Rep}_{\mathbb{F}[\![z]\!]}\,\mathbb{P}, \mathfrak{FMod}_{\mathbb{F}(\!(z)\!)[\pi_1(S,\bar{s})]})$, denote the category whose objects are tensor functors from $\mathrm{Rep}_{\mathbb{F}[\![z]\!]}\,\mathbb{P}$ to $\mathfrak{FMod}_{\mathbb{F}[\![z]\!][\pi_1(S,\bar{s})]}$, respectively to $\mathfrak{FMod}_{\mathbb{F}(\!(z)\!)[\pi_1(S,\bar{s})]}$, and whose morphisms are isomorphisms of functors. We define the (*dual*) *Tate functor* $\check{\mathcal{T}}_{-}$, respectively the *rational* (*dual*) *Tate functor* $\check{\mathcal{V}}_{-}$ as the functors

$$\begin{array}{rccl} \check{\mathcal{T}}_{-}\colon & Sht_{\mathbb{P}}^{\mathbb{D}}(S) & \longrightarrow & Funct^{\otimes}(\mathrm{Rep}_{\mathbb{F}[\![z]\!]}\,\mathbb{P},\ \mathfrak{FMod}_{\mathbb{F}[\![z]\!][\pi_1(S,\bar{s})]}) \\ & \underline{\mathcal{L}} & \longmapsto & \check{\mathcal{T}}_{\underline{\mathcal{L}}}\colon\ \rho \mapsto \check{T}_{\rho_*\underline{\mathcal{L}}}, \\ \check{\mathcal{V}}_{-}\colon & Sht_{\mathbb{P}}^{\mathbb{D}}(S) & \longrightarrow & Funct^{\otimes}(\mathrm{Rep}_{\mathbb{F}[\![z]\!]}\,\mathbb{P},\ \mathfrak{FMod}_{\mathbb{F}(\!(z)\!)[\pi_1(S,\bar{s})]}) \\ & \underline{\mathcal{L}} & \longmapsto & \check{\mathcal{V}}_{\underline{\mathcal{L}}}\colon\ \rho \mapsto \check{V}_{\rho_*\underline{\mathcal{L}}}. \end{array}$$

If $\underline{\mathcal{L}}$ is an étale local $\mathbb{P}$-shtuka then the composition of the tensor functor $\check{\mathcal{T}}_{\underline{\mathcal{L}}}$ followed by the forgetful functor $F\colon \mathfrak{FMod}_{\mathbb{F}[\![z]\!][\pi_1(S,\bar{s})]} \to \mathfrak{FMod}_{\mathbb{F}[\![z]\!]}$ is isomorphic to the forgetful fiber functor $\omega^\circ\colon\ \mathrm{Rep}_{\mathbb{F}[\![z]\!]}\,\mathbb{P} \to \mathfrak{FMod}_{\mathbb{F}[\![z]\!]}$ by Corollary 2.9. Indeed, the base change $\underline{\mathcal{L}}_{\bar{s}}$ of $\underline{\mathcal{L}}$ to $\bar{s} = \operatorname{Spec}\kappa(\bar{s})$ is isomorphic to $\underline{\mathbb{L}}_0 := \big((L^+\mathbb{P})_{\bar{s}}, 1\cdot\hat{\sigma}^*\big)$ and the functor $F \circ \check{\mathcal{T}}_{\underline{\mathbb{L}}_0}$ is isomorphic to ω°. Consider the full subcategory $Funct^{\otimes}_0(\mathrm{Rep}_{\mathbb{F}[\![z]\!]}\,\mathbb{P},\ \mathfrak{FMod}_{\mathbb{F}[\![z]\!][\pi_1(S,\bar{s})]})$ of $Funct^{\otimes}(\mathrm{Rep}_{\mathbb{F}[\![z]\!]}\,\mathbb{P},\ \mathfrak{FMod}_{\mathbb{F}[\![z]\!][\pi_1(S,\bar{s})]})$ consisting of those tensor functors $\mathcal{F}$ for which $F\circ\mathcal{F} \cong \omega^\circ$. Then Proposition 3.4 generalizes as follows.

Proposition 3.6. *Let $S \in \mathcal{N}ilp_{\mathbb{F}[\![\zeta]\!]}$ be connected. Then the functor $\check{\mathcal{T}}_{-}$ is an equivalence between the category of* étale *local $\mathbb{P}$-shtukas over S and the category $Funct^{\otimes}_0(\mathrm{Rep}_{\mathbb{F}[\![z]\!]}\,\mathbb{P},\ \mathfrak{FMod}_{\mathbb{F}[\![z]\!][\pi_1(S,\bar{s})]})$. The functor $\check{\mathcal{V}}_{-}$ from the category of* étale *local $\mathbb{P}$-shtukas over S with quasi-isogenies as morphisms to the category $Funct^{\otimes}(\mathrm{Rep}_{\mathbb{F}[\![z]\!]}\,\mathbb{P},\ \mathfrak{FMod}_{\mathbb{F}(\!(z)\!)[\pi_1(S,\bar{s})]})$ is fully faithful.*

Proof. To construct the functor which is quasi-inverse to $\check{\mathcal{T}}_{-}$ we fix a tensor functor $\mathcal{F}$ in $Funct^{\otimes}_0(\mathrm{Rep}_{\mathbb{F}[\![z]\!]}\,\mathbb{P},\ \mathfrak{FMod}_{\mathbb{F}[\![z]\!][\pi_1(S,\bar{s})]})$ and an isomorphism $F\circ\mathcal{F} \cong \omega^\circ$. Since $\mathbb{P} = \mathrm{Aut}^{\otimes}(\omega^\circ)$ by [Wed04, Corollary 5.20] and $\pi_1(S,\bar{s})$ acts as automorphisms of the fiber functor $F\circ\mathcal{F}$, the functor $\mathcal{F}$ corresponds to a representation $\pi\colon \pi_1(S,\bar{s}) \to \mathbb{P}(\mathbb{F}[\![z]\!])$. For each $m \in \mathbb{N}$ we let $S_m \to S$ be the finite étale Galois covering corresponding to the kernel of $\pi_1(S,\bar{s}) \xrightarrow{\ \pi\ } \mathbb{P}(\mathbb{F}[\![z]\!]) \xrightarrow{\ \mathrm{mod}\, z^m\ } \mathbb{P}_m\big(\mathbb{F}[\![z]\!]/(z^m)\big)$ where $\mathbb{P}_m := \mathbb{P}\times_{\mathbb{D}} \operatorname{Spec}\mathbb{F}[\![z]\!]/(z^m)$. Let $\widetilde{\mathcal{G}}_m$ be the trivial $\mathbb{P}_m$-torsor over $S_m \times_{\mathbb{F}} \operatorname{Spec}\mathbb{F}[\![z]\!]/(z^m)$ and equip it with the Frobenius $\hat{\tau} := \mathrm{id}\colon \hat{\sigma}^*\widetilde{\mathcal{G}}_m \xrightarrow{\ \sim\ } \widetilde{\mathcal{G}}_m$. Via the action of $\mathrm{Gal}(S_m/S)$ through π on $\widetilde{\mathcal{G}}_m$ the latter descends to a $\mathbb{P}_m$-torsor $\mathcal{G}_m$ over $S \times_{\mathbb{F}} \operatorname{Spec}\mathbb{F}[\![z]\!]/(z^m)$ and $\hat{\tau}$ descends to an isomorphism $\hat{\tau}\colon \hat{\sigma}^*\mathcal{G}_m \to \mathcal{G}_m$. This makes $\hat{\mathcal{G}} := \varprojlim \mathcal{G}_m$ into a formal $\hat{\mathbb{P}}$-torsor over S together with an isomorphism $\hat{\tau}\colon \hat{\sigma}^*\hat{\mathcal{G}} \xrightarrow{\ \sim\ } \hat{\mathcal{G}}$; see Definition 2.2. By Proposition 2.4 it corresponds to an $L^+\mathbb{P}$-torsor $\mathcal{L}_+$ together with an isomorphism $\hat{\tau}\colon \hat{\sigma}^*\mathcal{L}_+ \xrightarrow{\ \sim\ } \mathcal{L}_+$, that is, to the étale local $\mathbb{P}$-shtuka $\underline{\mathcal{L}} = (\mathcal{L}_+, \hat{\tau})$. It satisfies $\check{\mathcal{T}}_{\underline{\mathcal{L}}} \cong \mathcal{F}$ and this yields the quasi-inverse to $\check{\mathcal{T}}_{-}$.

To prove that $\check{\mathcal{V}}_{-}$ is fully faithful let $\underline{\mathcal{L}} = (\mathcal{L}_+, \hat{\tau})$ and $\underline{\mathcal{L}}' = (\mathcal{L}'_+, \hat{\tau}')$ be two étale local $\mathbb{P}$-shtukas over S and let $\delta\colon \check{\mathcal{V}}_{\underline{\mathcal{L}}} \xrightarrow{\ \sim\ } \check{\mathcal{V}}_{\underline{\mathcal{L}}'}$ be an isomorphism of tensor functors. We consider the following functor

$$\widehat{\mathcal{M}}_{-}\colon \mathscr{H}^1(\operatorname{Spec}\mathbb{F}, L^+\mathbb{P})(S) \to Funct^{\otimes}(\mathrm{Rep}_{\mathbb{F}[\![z]\!]}\,\mathbb{P}, \mathfrak{Mod}_{\mathcal{O}_S[\![z]\!]}),$$

which sends an $L^+\mathbb{P}$-torsor $\mathcal{L}_+$ to the tensor functor mapping the representation ρ to the $\mathcal{O}_S[\![z]\!]$-module $\rho_*\mathcal{L}_+$ from (3.1). By Proposition 3.4 the isomorphism $\delta|_\rho$ between $\check{\mathcal{V}}_{\underline{\mathcal{L}}}(\rho) = \check{V}_{\rho_*\underline{\mathcal{L}}}$ and $\check{\mathcal{V}}_{\underline{\mathcal{L}}'}(\rho) = \check{V}_{\rho_*\underline{\mathcal{L}}'}$ comes from a quasi-isogeny between $\rho_*\underline{\mathcal{L}} = \big(\widehat{\mathcal{M}}_{\mathcal{L}_+}(\rho), \rho_*\hat{\tau}\big)$ and $\rho_*\underline{\mathcal{L}}' = \big(\widehat{\mathcal{M}}_{\mathcal{L}'_+}(\rho), \rho_*\hat{\tau}'\big)$. Therefore the isomorphism δ induces an isomorphism $\widehat{\mathcal{M}}_{\mathcal{L}_+} \otimes_{\mathcal{O}_S[\![z]\!]} \mathcal{O}_S(\!(z)\!) \xrightarrow{\ \sim\ } \widehat{\mathcal{M}}_{\mathcal{L}'_+} \otimes_{\mathcal{O}_S[\![z]\!]} \mathcal{O}_S(\!(z)\!)$. Take an étale cover $S' \to S$ trivializing $\mathcal{L}_+$ and $\mathcal{L}'_+$ and fix trivializations $\mathcal{L}_+ \cong (L^+\mathbb{P})_{S'} \cong \mathcal{L}'_+$. Then we have

$$\mathrm{Isom}^{\otimes}\big(\widehat{\mathcal{M}}_{(\mathcal{L}_+)_{S'}} \otimes_{\mathcal{O}_{S'}[\![z]\!]} \mathcal{O}_{S'}(\!(z)\!)\,,\, \widehat{\mathcal{M}}_{(\mathcal{L}_+)_{S'}} \otimes_{\mathcal{O}_{S'}[\![z]\!]} \mathcal{O}_{S'}(\!(z)\!)\big) \;\cong\; \mathrm{Aut}^{\otimes}(\omega^\circ)\big(\mathcal{O}_{S'}(\!(z)\!)\big) \;=\; LP(S'),$$

because $\widehat{\mathcal{M}}_{(L^+\mathbb{P})_{S'}} = \omega^\circ \otimes_{\mathbb{F}[\![z]\!]} \mathcal{O}_{S'}(\!(z)\!)$ and $\mathbb{P} = \mathrm{Aut}^{\otimes}(\omega^\circ)$ by [Wed04, Corollary 5.20]. Therefore the isomorphism δ gives an isomorphism $h_{S'}\colon (LP)_{S'} \to (LP)_{S'}$. The morphism $h_{S'}$ inherits the descent datum coming from the fact that δ is defined over S, and hence it defines an isomorphism $h\colon \mathcal{L} \xrightarrow{\sim} \mathcal{L}'$, where $\mathcal{L}$ and $\mathcal{L}'$ denote the LP-torsors associated with $\mathcal{L}_+$ and $\mathcal{L}'_+$. One easily checks that h satisfies $\hat\tau' \circ \hat\sigma^* h = h \circ \hat\tau$ and gives a quasi-isogeny $h\colon \underline{\mathcal{L}} \to \underline{\mathcal{L}}'$. □

Remark 3.7. If $\mathbb{P}(\mathbb{F}[\![z]\!]) \subset P(\mathbb{F}(\!(z)\!))$ is not a maximal compact subgroup the functor $\check{\mathcal{V}}_-$ does not need to be an equivalence, not even onto the category of those tensor functors $\mathcal{F}\colon \mathrm{Rep}_{\mathbb{F}[\![z]\!]}\,\mathbb{P} \to \mathfrak{FMod}_{\mathbb{F}(\!(z)\!)[\pi_1(S,\bar s)]}$ for which $F \circ \mathcal{F} = \omega^\circ \otimes_{\mathbb{F}[\![z]\!]} \mathbb{F}(\!(z)\!)$. For example let $\mathbb{P}$ be the Iwahori subgroup of GL_2, that is, $\mathbb{P}(\mathbb{F}[\![z]\!]) = \{\, A \in \mathrm{GL}_2(\mathbb{F}[\![z]\!])\colon A \equiv \left(\begin{smallmatrix} * & * \\ 0 & * \end{smallmatrix}\right) \bmod z \,\}$. Let x be transcendental over $\mathbb{F}$ and let $S = \mathrm{Spec}\,\mathbb{F}(x)$. Set $G := \mathrm{Gal}\big(\mathbb{F}(x)^{\mathrm{sep}}/\mathbb{F}(x)\big) = \pi_1(S,\bar s)$ and consider a representation $\pi\colon G \to \mathrm{GL}_2(\mathbb{F}[\![z]\!])$ such that the residual representation $\overline{\pi}\colon G \to \mathrm{GL}_2(\mathbb{F})$ is irreducible. Then $\pi(G) \not\subset \mathbb{P}(\mathbb{F}[\![z]\!])$ and hence the tensor functor $\mathcal{F}\colon \mathrm{Rep}_{\mathbb{F}[\![z]\!]}\,\mathbb{P} \to \mathfrak{FMod}_{\mathbb{F}(\!(z)\!)[G]}$ given by $\rho \longmapsto \big[G \xrightarrow{\pi} \mathrm{GL}_2(\mathbb{F}[\![z]\!]) \hookrightarrow P\big(\mathbb{F}(\!(z)\!)\big) \xrightarrow{\rho} \mathrm{GL}\big(\omega^\circ(\rho)\big)\big(\mathbb{F}(\!(z)\!)\big)\big]$ cannot come from a local $\mathbb{P}$-shtuka over $\mathbb{F}(x)$.

Note that such a representation π exists. For example if $\varphi\colon \mathbb{F}_q[t] \to \mathrm{End}\,\mathbb{G}_{a,\mathbb{F}(x)}$ is a Drinfeld-$\mathbb{F}_q[t]$-module of rank 2 over $\mathbb{F}(x)$ without potential complex multiplication, then for almost all primes ν of $\mathbb{F}_q[t]$ the Galois representation $\pi_{\varphi,\nu}\colon G \to \mathrm{GL}_2(A_\nu)$ on the ν-adic Tate module of φ has this property by [PT06, Theorem A] or [PR09, Theorem 0.1]. For a concrete example let $\varphi_t = 1 - x\tau + \tau^2$ and $\nu = (t)$. Then $\varphi[t] = \{y \in \mathbb{F}(x)^{\mathrm{sep}}\colon y^{q^2} - xy^q + y = 0\}$. If $q = 2$ it is easy to see that $y^3 - xy + 1$ is irreducible in $\mathbb{F}(x)[y]$ and has splitting field of degree 6 over $\mathbb{F}(x)$. This implies that $\overline{\pi}_{\varphi,(t)}(G) = \mathrm{GL}_2(\mathbb{F}_2)$. The reason for the failure of $\check{\mathcal{V}}_-$ to be an equivalence of course lies in the fact that the Drinfeld-module φ does not carry a level structure over $\mathbb{F}(x)$ whereas any étale local $\mathbb{P}$-shtuka for the above Iwahori group $\mathbb{P}$ carries a $\Gamma_0(\nu)$-level structure.

4 The Rapoport-Zink Spaces for Local ℙ-Shtukas

Rapoport and Zink constructed a moduli space for p-divisible groups together with a quasi-isogeny to a fixed one (and with some extra structure such as a polarization, endomorphisms, or a level structure). They proved that this moduli space is pro-representable by a formal scheme locally formally of finite type over $\mathbb{Z}_p$.

In this chapter we assume that $\mathbb{P}$ is a parahoric group scheme over $\mathbb{D}$ in the sense of [BT72, Définition 5.2.6] and [HR03]. We already mentioned that local $\mathbb{P}$-shtukas behave analogously to p-divisible groups. It turns out that this analogy is not perfect, unless we restrict to "bounded" local $\mathbb{P}$-shtukas as the analogous objects corresponding to p-divisible groups. More precisely we bound the *Hodge polygon* of a local $\mathbb{P}$-shtuka $(\mathcal{L}_+, \hat\tau)$, that is, the relative position of $\hat\sigma^*\mathcal{L}_+$ and $\mathcal{L}_+$ under the isomorphism $\hat\tau$. We will show that Rapoport-Zink spaces for bounded local $\mathbb{P}$-shtukas are formal schemes locally formally of finite type over $\mathrm{Spf}\,\mathbb{F}[\![\zeta]\!]$. When $\mathbb{P}$ is constant, i.e. $\mathbb{P} := G_0 \times_{\mathbb{F}} \mathbb{D}$ for a reductive group G_0 over $\mathbb{F}$ this was proved in [HV11, Theorem 6.3]. In the non-constant case for a parahoric group $\mathbb{P}$ over $\mathbb{D}$ we will give an axiomatic definition of the boundedness condition in Section 4.2. We start with the unbounded situation.

4.1 Unbounded Rapoport-Zink Spaces

For a scheme S in $\mathcal{N}ilp_{\mathbb{F}[\![\zeta]\!]}$ let $\bar S$ denote the closed subscheme $\mathrm{V}_S(\zeta) \subseteq S$. On the other hand for a scheme $\bar T$ over $\mathbb{F}$ we set $\widehat{T} := \bar T \,\widehat{\times}_{\mathrm{Spec}\,\mathbb{F}}\, \mathrm{Spf}\,\mathbb{F}[\![\zeta]\!]$. Then $\widehat{T}$ is a ζ-adic formal scheme with underlying topological space $\bar T = \mathrm{V}_{\widehat{T}}(\zeta)$.

Definition 4.1. With a given local $\mathbb{P}$-shtuka $\underline{\mathcal{L}}_0$ over an $\mathbb{F}$-scheme $\bar{T}$ we associate the functor

$$\begin{aligned}\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}\colon \mathcal{N}ilp_{\widehat{T}} &\longrightarrow \mathcal{S}ets\\ S &\longmapsto \big\{\text{Isomorphism classes of } (\underline{\mathcal{L}},\bar{\delta})\colon \text{ where } \underline{\mathcal{L}} \text{ is a local } \mathbb{P}\text{-shtuka}\\ &\qquad\quad \text{over } S \text{ and } \bar{\delta}\colon \underline{\mathcal{L}}_{\bar{S}} \to \underline{\mathcal{L}}_{0,\bar{S}} \text{ is a quasi-isogeny over } \bar{S}\big\}.\end{aligned}$$

Here we say that $(\underline{\mathcal{L}},\bar{\delta})$ and $(\underline{\mathcal{L}}',\bar{\delta}')$ are isomorphic if $\bar{\delta}^{-1}\circ\bar{\delta}'$ lifts to an isomorphism $\underline{\mathcal{L}}'\to\underline{\mathcal{L}}$. The group $\mathrm{QIsog}_{\bar{T}}(\underline{\mathcal{L}}_0)$ of quasi-isogenies of $\underline{\mathcal{L}}_0$ acts on the functor $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}$ via $g\colon(\underline{\mathcal{L}},\bar{\delta})\mapsto(\underline{\mathcal{L}},g\circ\bar{\delta})$ for $g\in\mathrm{QIsog}_{\bar{T}}(\underline{\mathcal{L}}_0)$. We will show that $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}$ is representable by an ind-scheme which we call an *unbounded Rapoport-Zink space for local* $\mathbb{P}$*-shtukas.*

Remark 4.2. Note that by rigidity of quasi-isogenies (Proposition 2.11) the functor $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}$ is naturally isomorphic to the functor

$$\begin{aligned}S \;\mapsto\; &\big\{\text{Isomorphism classes of } (\underline{\mathcal{L}},\delta)\colon \text{ where } \underline{\mathcal{L}} \text{ is a local } \mathbb{P}\text{-shtuka}\\ &\qquad\quad \text{over } S \text{ and } \delta\colon \underline{\mathcal{L}}_{S} \to \underline{\mathcal{L}}_{0,S} \text{ is a quasi-isogeny over } S\big\}.\end{aligned}$$

This also shows that $\mathrm{id}_{\underline{\mathcal{L}}}$ is the only automorphism of $(\underline{\mathcal{L}},\delta)$, and for this reason we do not need to consider $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}$ as a stack.

Remark 4.3. In order to show that $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}$ is representable by an ind-scheme we recall the definition of the affine flag variety $\mathcal{F}\ell_{\mathbb{P}}$. It is defined to be the *fpqc*-sheaf associated with the presheaf

$$R \longmapsto LP(R)/L^+\mathbb{P}(R) \;=\; \mathbb{P}\left(R(\!(z)\!)\right)/\mathbb{P}\left(R[\![z]\!]\right)$$

on the category of $\mathbb{F}$-algebras. Pappas and Rapoport [PR08, Theorem 1.4] show that $\mathcal{F}\ell_{\mathbb{P}}$ is represented by an ind-scheme which is ind-quasi-projective over $\mathbb{F}$, and hence ind-separated and of ind-finite type over $\mathbb{F}$. Moreover, they show that the quotient morphism $LP\to\mathcal{F}\ell_{\mathbb{P}}$ admits sections locally for the étale topology. They proceed as follows. When $\mathbb{P}=\mathrm{SL}_{r,\mathbb{D}}$, the *fpqc*-sheaf $\mathcal{F}\ell_{\mathbb{P}}$ is called the *affine Grassmanian.* It is an inductive limit of projective schemes over $\mathbb{F}$, that is, ind-projective over $\mathbb{F}$; see [BD, Theorem 4.5.1] or [Fal03, NP01]. By Proposition 2.13(a) there is a faithful representation $\mathbb{P}\hookrightarrow\mathrm{SL}_r$ with quasi-affine quotient. Pappas and Rapoport show in the proof of [PR08, Theorem 1.4] that $\mathcal{F}\ell_{\mathbb{P}}\to\mathcal{F}\ell_{\mathrm{SL}_r}$ is a locally closed embedding, and moreover, if $\mathrm{SL}_r/\mathbb{P}$ is affine, then $\mathcal{F}\ell_{\mathbb{P}}\to\mathcal{F}\ell_{\mathrm{SL}_r}$ is even a closed embedding. In particular we see from Proposition 2.13(b) that $\mathcal{F}\ell_{\mathbb{P}}$ is ind-projective if $\mathbb{P}$ is reductive.

Let us view the formal scheme $\widehat{T}=\bar{T}\,\widehat{\times}_{\mathbb{F}}\,\mathrm{Spf}\,\mathbb{F}[\![\zeta]\!]$ as the ind-scheme $\varinjlim \bar{T}\times_{\mathbb{F}}\mathrm{Spec}\,\mathbb{F}[\zeta]/(\zeta^m)$. We may form the fiber product $\widehat{\mathcal{F}\ell}_{\mathbb{P},\bar{T}} := \mathcal{F}\ell_{\mathbb{P}}\,\widehat{\times}_{\mathbb{F}}\,\widehat{T}$ in the category of ind-schemes (see [BD, 7.11.1]). Note that this fiber product can be either viewed as the restriction of the sheaf $\mathcal{F}\ell_{\mathbb{P}}$ to the *fpqc*-site of schemes in $\mathcal{N}ilp_{\widehat{T}}$ or also as the formal completion of $\mathcal{F}\ell_{\mathbb{P}}\times_{\mathbb{F}}\bar{T}\times_{\mathbb{F}}\mathrm{Spec}\,\mathbb{F}[\![\zeta]\!]$ along the special fiber $\mathrm{V}(\zeta)$.

Theorem 4.4. *The functor $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}$ from Definition 4.1 is represented by an ind-scheme, ind-quasi-projective over $\widehat{T}=\bar{T}\,\widehat{\times}_{\mathbb{F}}\,\mathrm{Spf}\,\mathbb{F}[\![\zeta]\!]$, hence ind-separated and of ind-finite type over $\widehat{T}$.*

(a) If $\underline{\mathcal{L}}_0$ is trivialized by an isomorphism $\alpha\colon\underline{\mathcal{L}}_0\xrightarrow{\sim}\big((L^+\mathbb{P})_{\bar{T}},b\hat{\sigma}^*\big)$ with $b\in LP(\bar{T})$ then $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}$ is represented by the ind-scheme $\widehat{\mathcal{F}\ell}_{\mathbb{P},\bar{T}}:=\mathcal{F}\ell_{\mathbb{P}}\,\widehat{\times}_{\mathbb{F}}\,\widehat{T}$.

(b) If $\mathbb{P}$ is reductive (or more generally if $\mathcal{F}\ell_{\mathbb{P}}$ is ind-projective over $\mathbb{F}$) then $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}$ is ind-projective over $\widehat{T}$.

Proof. We first prove statement (a). We regard $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}$ in the equivalent form mentioned in Remark 4.2. Consider a pair $(\underline{\mathcal{L}}, \delta) = ((\mathcal{L}_+, \hat{\tau}), \delta) \in \underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}(S)$. Choose an étale covering $S' \to S$ which trivializes $\underline{\mathcal{L}}$, then the quasi-isogeny $\alpha_{S'} \circ \delta$ is given by an element $g' \in LP(S')$. The image of the element $g' \in LP(S')$ in $\widehat{\mathcal{F}\ell}_{\mathbb{P},\bar{T}}(S')$ is independent of the choice of the trivialization, and since $(\underline{\mathcal{L}}, \delta)$ is defined over S, it descends to a point $x \in \widehat{\mathcal{F}\ell}_{\mathbb{P},\bar{T}}(S)$. Note in particular that $\hat{\tau}_{S'}$ is determined by b and g' through the diagram

$$\begin{tikzcd}
\hat{\sigma}^*(LP)_{S'} \arrow[r, "\hat{\tau}_{S'}"] \arrow[d, "\hat{\sigma}^*(g')"'] & (LP)_{S'} \arrow[d, "g'"] \\
\hat{\sigma}^*(LP)_{S'} \arrow[r, "b"] & (LP)_{S'}.
\end{tikzcd}$$

Conversely let $x \in \widehat{\mathcal{F}\ell}_{\mathbb{P},\bar{T}}(S)$ for a scheme S in $\mathcal{N}ilp_{\widehat{T}}$. The projection morphism $LP \to \mathcal{F}\ell_{\mathbb{P}}$ admits local sections for the étale topology by [PR08, Theorem 1.4]. Consequently there is an étale covering $S' \to S$ such that x is represented by an element $g' \in LP(S')$. We set $S'' := S' \times_S S'$ and define $(\mathcal{L}'_+, \hat{\tau}', \delta')$ over S' as follows. Let $\mathcal{L}'_+ := (L^+\mathbb{P})_{S'}$, let the quasi-isogeny $\delta'\colon (\mathcal{L}'_+, \hat{\tau}') \to \big((L^+\mathbb{P})_{S'}, b\hat{\sigma}^*\big)$ be given by $y \mapsto g'y$, and the Frobenius by $\hat{\tau}' := (g')^{-1} b \hat{\sigma}^*(g')\hat{\sigma}^*$. We descend $(\mathcal{L}'_+, \hat{\tau}', \delta')$ to S. For an S-scheme Y let $Y' = Y \times_S S'$ and $Y'' = Y' \times_Y Y' = Y \times_S S''$, and let $p_i\colon Y'' \to Y'$ be the projection onto the i-th factor. Since g' comes from an element $x \in \widehat{\mathcal{F}\ell}_{\mathbb{P},\bar{T}}(S)$ there is an $h \in L^+\mathbb{P}(S'')$ with $p_1^*(g') = p_2^*(g') \cdot h$. Consider the *fpqc*-sheaf $\mathcal{L}_+$ on S whose sections over an S-scheme Y are given by

$$\mathcal{L}_+(Y) \;:=\; \big\{\, y' \in L^+\mathbb{P}(Y') : \;\; p_1^*(y') = h^{-1} \cdot p_2^*(y') \text{ in } L^+\mathbb{P}(Y'') \,\big\}$$

on which $L^+\mathbb{P}(Y)$ acts by right multiplication. Then $\underline{\mathcal{L}}$ is a $L^+\mathbb{P}$-torsor on S because over $Y = S'$ there is a trivialization

$$(L^+\mathbb{P})_{S'} \xrightarrow{\sim} (\mathcal{L}_+)_{S'}\,, \qquad f \mapsto h\, p_1^*(f) \;\in\; (L^+\mathbb{P})(S'')$$

due to the cocycle condition for h. Moreover, $\hat{\tau}'$ descends to an isomorphism

$$\hat{\tau}\colon \hat{\sigma}^*\mathcal{L}(Y) \xrightarrow{\sim} \mathcal{L}(Y)\,,\; \hat{\sigma}^*(y') \mapsto (g')^{-1} b \hat{\sigma}^*(g') \hat{\sigma}^*(y')$$

making $(\mathcal{L}_+, \hat{\tau})$ into a local $\mathbb{P}$-shtuka over S. Also δ' descends to a quasi-isogeny of local $\mathbb{P}$-shtukas

$$\begin{array}{rcl}
\delta\colon \mathcal{L}(Y) & \xrightarrow{\sim} & LP(Y) = \big\{\, f' \in LP(Y') : \; p_1^*(f') = p_2^*(f') \text{ in } LP(Y'') \,\big\}\,, \\
y' & \mapsto & g'y'\,.
\end{array}$$

Note that this is well defined. Namely, if g' is replaced by $\tilde{g}'$ with $u' = (\tilde{g}')^{-1} g' \in L^+\mathbb{P}(S')$ then left multiplication with u' defines an isomorphism

$$\big((L^+\mathbb{P})_{S'}, (g')^{-1} b \hat{\sigma}^*(g')\hat{\sigma}^*, g'\big) \xrightarrow{\sim} \big((L^+\mathbb{P})_{S'}, (\tilde{g}')^{-1} b \hat{\sigma}^*(\tilde{g}')\hat{\sigma}^*, \tilde{g}'\big).$$

Also $\tilde{h} = p_2^*(u')\, h\, p_1^*(u')^{-1}$ and hence left multiplication with u' descends to an isomorphism $\underline{\mathcal{L}} \xrightarrow{\sim} \underline{\widetilde{\mathcal{L}}}$ over S. This establishes (a).

To prove the first assertion of the theorem we may choose an étale covering $\bar{T}' \to \bar{T}$ and a trivialization $\alpha\colon \underline{\mathcal{L}}_{0,\bar{T}'} \xrightarrow{\sim} \big((L^+\mathbb{P})_{\bar{T}'}, b\hat{\sigma}^*\big)$ over $\bar{T}'$. We set $\widehat{T}' := \bar{T}' \,\widehat{\times}_{\mathbb{F}} \operatorname{Spf} \mathbb{F}[\![\zeta]\!]$ and $\widehat{T}'' := \widehat{T}' \,\widehat{\times}_{\widehat{T}} \widehat{T}'$, and let $pr_i\colon \widehat{T}'' \to \widehat{T}'$ be the projection onto the i-th factor. Then (a) yields an isomorphism $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0} \,\widehat{\times}_{\widehat{T}} \widehat{T}' \xrightarrow{\sim}$

$\widehat{\mathcal{F}\ell}_{\mathbb{P},\bar T'}$, $\delta \mapsto \alpha_{S'} \circ \delta = g' \cdot L^+\mathbb{P}(S')$. Note that $\widehat{\mathcal{F}\ell}_{\mathbb{P},\bar T'}$ is an ind-scheme which is ind-quasi-projective over $\widehat T'$ by Remark 4.3. Over $\widehat T''$ there is an isomorphism

$$\begin{tikzcd}
pr_1^*(\widehat{\mathcal{F}\ell}_{\mathbb{P},\bar T'}) \arrow[r, "\cong"] & \underline{\mathcal{M}}_{\underline{\mathcal{L}}_0} \hat\times_{\widehat T} \widehat T'' \arrow[r, "\cong"] & pr_2^*(\widehat{\mathcal{F}\ell}_{\mathbb{P},\bar T'}) \\
g' \cdot L^+\mathbb{P}(S') \arrow[r, mapsto] & pr_1^*\alpha_{S'}^{-1} \circ g' \arrow[r, mapsto] & pr_2^*\alpha_{S'} \circ pr_1^*\alpha_{S'}^{-1} \circ g' \cdot L^+\mathbb{P}(S')
\end{tikzcd} \tag{4.1}$$

It is given by left multiplication on $\widehat{\mathcal{F}\ell}_{\mathbb{P},\bar T''} = pr_i^*(\widehat{\mathcal{F}\ell}_{\mathbb{P},\bar T'})$ with the element $pr_2^*\alpha_{S'} \circ pr_1^*\alpha_{S'}^{-1} \in L^+\mathbb{P}(\widehat T'')$.

We write $\mathcal{F}\ell_{\mathbb{P}} = \varinjlim \mathcal{F}\ell_{\mathbb{P}}^{(N)}$ for quasi-projective $\mathbb{F}$-schemes $\mathcal{F}\ell_{\mathbb{P}}^{(N)}$. There is a line bundle $\mathcal{F}$ on $\mathcal{F}\ell_{\mathbb{P}}$ which is "ample" in the sense that its restriction to any $\mathcal{F}\ell_{\mathbb{P}}^{(N)}$ is ample, and which is equivariant for the $L^+\mathbb{P}$-action by left multiplication, i.e. "$L^+\mathbb{P}$-linearized" in the sense of [MFK94, Definition 1.6]. For example one can take a faithful representation $\mathbb{P} \hookrightarrow \mathrm{SL}_{r,\mathbb{D}}$ with quasi-affine quotient (Proposition 2.13(a)), take the fundamental line bundles $\mathcal{L}_i$ on $\mathcal{F}\ell_{\mathrm{SL}_r}$ from [Fal03, p. 46], take $\mathcal{F}$ as the pullback of a tensor product of strictly positive powers of the $\mathcal{L}_i$, see [Fal03, p. 54], and take the $\mathcal{F}\ell_{\mathbb{P}}^{(N)}$ as the preimages in $\mathcal{F}\ell_{\mathbb{P}}$ of the Schubert varieties in $\mathcal{F}\ell_{\mathrm{SL}_r}$. Then (4.1) defines a descent datum on the pair $(\mathcal{F}\ell_{\mathbb{P}}^{(N)} \times_{\mathbb{F}} \bar T' \times_{\mathbb{F}} \mathbb{F}[\![\zeta]\!]/(\zeta^m), \mathcal{F})$ for all N and m. This descent datum is effective by [SGA 1, VIII, Proposition 7.8], see also [BLR90, §6.2, Theorem 7]. Therefore there is a quasi-projective scheme $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}^{(N,m)}$ over $\bar T \times_{\mathbb{F}} \mathbb{F}[\![\zeta]\!]/(\zeta^m)$ with $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}^{(N,m)} \times_{\bar T} \bar T' \cong \mathcal{F}\ell_{\mathbb{P}}^{(N)} \times_{\mathbb{F}} \bar T' \times_{\mathbb{F}} \mathbb{F}[\![\zeta]\!]/(\zeta^m)$. It follows that $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0} = \varinjlim_{N,m} \underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}^{(N,m)}$ is an ind-quasi-projective ind-scheme over $\hat T$.

(b) If $\mathbb{P}$ is reductive then there is a representation $\mathbb{P} \hookrightarrow \mathrm{SL}_{r,\mathbb{D}}$ with affine quotient by Proposition 2.13(b) and then $\mathcal{F}\ell_{\mathbb{P}}^{(N)}$ is projective over $\mathbb{F}$ by Remark 4.3. Therefore also $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}^{(N,m)}$ is projective over $\bar T \times_{\mathbb{F}} \mathbb{F}[\![\zeta]\!]/(\zeta^m)$, and hence $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}$ is ind-projective over $\widehat T$. □

4.2 Bounded Local $\mathbb{P}$-Shtukas

Definition 4.5. We define a *bound* to be a closed ind-subscheme $\hat Z$ of $\widehat{\mathcal{F}\ell}_{\mathbb{P}} := \mathcal{F}\ell_{\mathbb{P}} \hat\times_{\mathbb{F}} \operatorname{Spf} \mathbb{F}[\![\zeta]\!]$ which is stable under the left $L^+\mathbb{P}$-action, such that $Z := \hat Z \hat\times_{\operatorname{Spf} \mathbb{F}[\![\zeta]\!]} \operatorname{Spec} \mathbb{F}$ is a quasi-compact subscheme of $\mathcal{F}\ell_{\mathbb{P}}$.

Let $\mathcal{L}_+$ and $\mathcal{L}'_+$ be $L^+\mathbb{P}$-torsors over a scheme S in $\mathcal{N}ilp_{\mathbb{F}[\![\zeta]\!]}$ and let $\delta\colon \mathcal{L} \xrightarrow{\sim} \mathcal{L}'$ be an isomorphism of the associated LP-torsors. We consider an étale covering $S' \to S$ over which trivializations $\alpha : \mathcal{L}_+ \xrightarrow{\sim} (L^+\mathbb{P})_{S'}$ and $\alpha' : \mathcal{L}'_+ \xrightarrow{\sim} (L^+\mathbb{P})_{S'}$ exist. Then the automorphism $\alpha' \circ \delta \circ \alpha^{-1}$ of $(LP)_{S'}$ corresponds to a morphism $S' \to LP \hat\times_{\mathbb{F}} \operatorname{Spf} \mathbb{F}[\![\zeta]\!]$. We say that δ is *bounded by* $\hat Z$ if for (some) any such trivialization the induced morphism $S' \to LP \hat\times_{\mathbb{F}} \operatorname{Spf} \mathbb{F}[\![\zeta]\!] \to \widehat{\mathcal{F}\ell}_{\mathbb{P}}$ factors through $\hat Z$. By the $L^+\mathbb{P}$-invariance of $\hat Z$ the definition is independent of the trivializations. Furthermore we say that a local $\mathbb{P}$-shtuka $(\mathcal{L}_+, \hat\tau)$ is *bounded by* $\hat Z$ if the isomorphism $\hat\tau$ is bounded by $\hat Z$.

In Examples 4.8 and 4.9 below we discuss the motivation for this definition and the relation to other boundedness conditions like in [HV11].

Remark 4.6. Let the ind-scheme structure on $\mathcal{F}\ell_{\mathbb{P}}$ be given as $\mathcal{F}\ell_{\mathbb{P}} = \varinjlim \mathcal{F}\ell_{\mathbb{P}}^{(m)}$ and the one on $\widehat{\mathcal{F}\ell}_{\mathbb{P}}$ as $\widehat{\mathcal{F}\ell}_{\mathbb{P}} = \varinjlim \mathcal{F}\ell_{\mathbb{P}}^{(m)} \times_{\mathbb{F}} \operatorname{Spec} \mathbb{F}[\![\zeta]\!]/(\zeta^m)$. Let $\hat Z^{(m)} := \hat Z \times_{\widehat{\mathcal{F}\ell}_{\mathbb{P}}} (\mathcal{F}\ell_{\mathbb{P}}^{(m)} \times_{\mathbb{F}} \operatorname{Spec} \mathbb{F}[\![\zeta]\!]/(\zeta^m))$. Then $\hat Z = \varinjlim \hat Z^{(m)}$ and $\hat Z_{\mathrm{red}} := \varinjlim \hat Z_{\mathrm{red}}^{(m)} = \varinjlim (\hat Z \hat\times_{\operatorname{Spf} \mathbb{F}[\![\zeta]\!]} \operatorname{Spec} \mathbb{F} \times_{\mathcal{F}\ell_{\mathbb{P}}} \mathcal{F}\ell_{\mathbb{P}}^{(m)})_{\mathrm{red}} = Z_{\mathrm{red}}$ is a scheme. This means that $\hat Z$ is a "reasonable formal scheme" over $\operatorname{Spf} \mathbb{F}[\![\zeta]\!]$ in the sense of [BD, 7.11.1 and 7.12.17], and hence a formal scheme in the sense of [EGA, $\mathrm{I}_{\mathrm{new}}$].

Proposition 4.7. *Let $\mathcal{L}_+$ and $\mathcal{L}'_+$ be $L^+\mathbb{P}$-torsors over a scheme $S \in \mathcal{N}ilp_{\mathbb{F}[\![\zeta]\!]}$ and let $\delta\colon \mathcal{L} \xrightarrow{\sim} \mathcal{L}'$ be an isomorphism of the associated LP-torsors. Let $\hat{Z}$ be a bound. Then the condition that δ is bounded by $\hat{Z}$ is represented by a closed subscheme of S.*

Proof. As in Definition 4.5 consider trivializations of $\mathcal{L}_+$ and $\mathcal{L}'_+$ over an étale covering $S' \to S$ and the induced morphism $S' \to LP \,\widehat{\times}_{\mathbb{F}} \operatorname{Spf} \mathbb{F}[\![\zeta]\!] \to \widehat{\mathcal{F}\ell}_{\mathbb{P}}$. By the $L^+\mathbb{P}$-invariance of $\hat{Z}$ the closed subscheme $S' \,\widehat{\times}_{\widehat{\mathcal{F}\ell}_{\mathbb{P}}} \hat{Z}$ of S' descends to a closed subscheme of S which represents the boundedness of δ by $\hat{Z}$. □

Example 4.8. (a) Let P be the generic fiber of $\mathbb{P}$ over $\operatorname{Spec} \mathbb{F}(\!(z)\!)$ and consider the base change P_L of P to $L = \mathbb{F}^{\mathrm{alg}}(\!(z)\!)$. Let A be a maximal split torus in P_L and let T be its centralizer. Since $\mathbb{F}^{\mathrm{alg}}$ is algebraically closed, P_L is quasi-split and so T is a maximal torus in P_L. Let $N = N(T)$ be the normalizer of T and let $\mathcal{T}^0$ be the identity component of the Néron model of T over $\mathcal{O}_L$.

The *Iwahori-Weyl group* associated with A is the quotient group $\widetilde{W} = N(L)/\mathcal{T}^0(\mathcal{O}_L)$. It is an extension of the finite Weyl group $W_0 = N(L)/T(L)$ by the coinvariants $X_*(T)_I$ under $I = \operatorname{Gal}(L^{\mathrm{sep}}/L)$:

$$0 \to X_*(T)_I \to \widetilde{W} \to W_0 \to 1.$$

By [HR03, Proposition 8] there is a bijection

$$L^+\mathbb{P}(\mathbb{F}^{\mathrm{alg}})\backslash LP(\mathbb{F}^{\mathrm{alg}})/L^+\mathbb{P}(\mathbb{F}^{\mathrm{alg}}) \xrightarrow{\sim} \widetilde{W}^{\mathbb{P}}\backslash\widetilde{W}/\widetilde{W}^{\mathbb{P}} \tag{4.2}$$

where $\widetilde{W}^{\mathbb{P}} := (N(L) \cap \mathbb{P}(\mathcal{O}_L))/\mathcal{T}^0(\mathcal{O}_L)$.

The *Schubert variety* $\mathcal{S}(\omega)$ associated with $\omega \in \widetilde{W}^{\mathbb{P}}\backslash\widetilde{W}/\widetilde{W}^{\mathbb{P}}$ is the ind-scheme theoretic closure of the $L^+\mathbb{P}$-orbit of ω in $\mathcal{F}\ell_{\mathbb{P}}$. It is a reduced projective variety over $\mathbb{F}$. For further details see [PR08] and [Ric10]. When $\hat{Z} = \mathcal{S}(\omega) \,\widehat{\times}_{\mathbb{F}} \operatorname{Spf} \mathbb{F}[\![\zeta]\!]$ for a Schubert variety $\mathcal{S}(\omega)$, with $\omega \in \widetilde{W}$, we say that δ is *bounded by* ω.

(b) In [HV11] Viehmann and the second author considered the case where $\mathbb{P} = G \times_{\mathbb{F}} \mathbb{D}$ for a split connected reductive group G over $\mathbb{F}$. In this case $\widetilde{W}^{\mathbb{P}} = W_0$ and $\widetilde{W}^{\mathbb{P}}\backslash\widetilde{W}/\widetilde{W}^{\mathbb{P}} = X_*(T)$. If $\mu \in X_*(T)$ one could consider the bound $\hat{Z} := \mathcal{S}(\mu) \,\widehat{\times}_{\mathbb{F}} \operatorname{Spf} \mathbb{F}[\![\zeta]\!]$ as in (a) above.

However, we instead fixed a Borel subgroup B of G and its opposite Borel $\overline{B}$. We considered a generating system Λ of the monoid of dominant weights $X^*(T)_{\mathrm{dom}}$, and for all $\lambda \in \Lambda$ the Weyl module $V_\lambda := \big(\operatorname{Ind}_{\overline{B}}^{G}(-\lambda)_{\mathrm{dom}}\big)^\vee$. For the representation $\rho_\lambda\colon G \to \operatorname{GL}(V_\lambda)$ we considered the sheaves of $\mathcal{O}_S[\![z]\!]$-modules $\rho_{\lambda*}\mathcal{L}_+$ and $\rho_{\lambda*}\mathcal{L}'_+$ associated with the $L^+\mathbb{P}$-torsors $\mathcal{L}_+$ and $\mathcal{L}'_+$ over S. (For the definition of $\mathcal{O}_S[\![z]\!]$ see Section 3.) The isomorphism δ of the associated LP-torsors corresponds to an isomorphism $\rho_{\lambda*}\delta\colon \rho_{\lambda*}\mathcal{L}_+ \otimes_{\mathcal{O}_S[\![z]\!]} \mathcal{O}_S(\!(z)\!) \xrightarrow{\sim} \rho_{\lambda*}\mathcal{L}'_+ \otimes_{\mathcal{O}_S[\![z]\!]} \mathcal{O}_S(\!(z)\!)$. We said in [HV11, Definition 3.5] that "δ is bounded by $(\tilde{z}, \mu)$", where $\tilde{z} = z$ or $\tilde{z} = z - \zeta$, if for all $\lambda \in \Lambda$

$$\rho_{\lambda*}\delta\,(\rho_{\lambda*}\mathcal{L}_+) \;\subset\; \tilde{z}^{\,-\langle\,(-\lambda)_{\mathrm{dom}},\mu\rangle}\,(\rho_{\lambda*}\mathcal{L}'_+),$$

and if for all geometric points $\bar{s}$ of S the image of the isomorphism $\delta_{\bar{s}}$ at $\bar{s}$ under the isomorphism (4.2) has the same image in $\pi_1(G)$ than μ.

In terms of Definition 4.5 this can be described as follows. Consider the universal matrix $M \in L\operatorname{GL}(V_\lambda)(S)$ over the ind-scheme $S_\lambda := L\operatorname{GL}(V_\lambda) \,\widehat{\times}_{\mathbb{F}} \operatorname{Spf} \mathbb{F}[\![\zeta]\!]$. Let $\overline{S}_\lambda$ be the closed ind-subscheme of S_λ defined by the condition that the matrix $\tilde{z}^{\langle(-\lambda)_{\mathrm{dom}},\mu\rangle}M$ has entries in $\mathcal{O}_{S_\lambda}[\![z]\!]$, and let

$$\hat{Z}_\lambda \;:=\; \overline{S}_\lambda/(L^+\operatorname{GL}(V_\lambda) \,\widehat{\times}_{\mathbb{F}} \operatorname{Spf} \mathbb{F}[\![\zeta]\!]) \;\subset\; \widehat{\mathcal{F}\ell}_{\operatorname{GL}(V_\lambda)}\,.$$

Let $(\widehat{\mathcal{F}\ell}_{\mathbb{P}})_{[\mu]}$ be the connected component of $\widehat{\mathcal{F}\ell}_{\mathbb{P}}$ corresponding to $[\mu] \in \pi_1(G)$ under the isomorphism $\pi_0(\widehat{\mathcal{F}\ell}_{\mathbb{P}}) \cong \pi_1(G)$; see [BD, Proposition 4.5.4] or [PR08, Theorem 0.1]. Write $\Lambda = \{\lambda_1, \ldots, \lambda_m\}$ and for each λ_i consider the morphism $\rho_{\lambda_i *}\colon (\widehat{\mathcal{F}\ell}_{\mathbb{P}})_{[\mu]} \to \widehat{\mathcal{F}\ell}_{\mathrm{GL}(V_{\lambda_i})}$ induced from ρ_{λ_i}. Then the base change $\hat{Z}$ of the closed ind-subscheme $\hat{Z}_{\lambda_1} \widehat{\times}_{\operatorname{Spf}\mathbb{F}[\![\zeta]\!]} \ldots \widehat{\times}_{\operatorname{Spf}\mathbb{F}[\![\zeta]\!]} \hat{Z}_{\lambda_m}$ under the morphism $\prod_i \rho_{\lambda_i *}\colon (\widehat{\mathcal{F}\ell}_{\mathbb{P}})_{[\mu]} \to \widehat{\mathcal{F}\ell}_{\mathrm{GL}(V_{\lambda_1})} \widehat{\times}_{\mathbb{F}[\![\zeta]\!]} \ldots \widehat{\times}_{\mathbb{F}[\![\zeta]\!]} \widehat{\mathcal{F}\ell}_{\mathrm{GL}(V_{\lambda_m})}$ is the bound representing the "boundedness by $(\tilde{z},\mu)$" from [HV11, Definition 3.5].

Instead of the Weyl modules V_λ one could of course also work with other representations of G. If for example one takes the induced modules $\mathrm{Ind}_B^G(-\lambda)_{\mathrm{dom}}$, or tilting modules, one obtains different ind-subschemes $\hat{Z}$, but the underlying reduced ind-subschemes of these $\hat{Z}$ still coincide. If $\tilde{z} = z$, this reduced ind-subscheme equals the Schubert variety $\mathcal{S}(\mu) \widehat{\times}_{\mathbb{F}} \operatorname{Spec}\mathbb{F}[\![\zeta]\!]$. This already indicates that it is reasonable to consider boundedness in terms of closed ind-subschemes of $\widehat{\mathcal{F}\ell}_{\mathbb{P}}$. Note that here also for $\tilde{z} = z - \zeta$ the bound $\hat{Z}$ only depends on μ and the class of modules considered (Weyl modules, etc.). This is no longer true if $\mathbb{P}$ is parahoric as one sees from the next example.

Example 4.9. Assume that $\operatorname{char}\mathbb{F}_q \neq 2$. Set $K := \mathbb{F}_q(\!(z)\!)$. Let $E := K[y]/(y^2 - z)$ be the ramified quadratic field extension with $y^2 = z$. Let T be the one dimensional torus $\ker(N_{E/K}\colon \operatorname{Res}_{E/K}\mathbb{G}_m \to \mathbb{G}_m)$. Explicitly $T = \operatorname{Spec} K[a,b]/(a^2 - b^2 z - 1)$, with the multiplication $(a,b)*(c,d) = (ac + bdz, ad + bc)$. Sending $a \mapsto \frac{1}{2}(t + t^{-1})$ and $b \mapsto \frac{1}{2y}(t^{-1} - t)$ defines an isomorphism $\mathbb{G}_{m,E} = \operatorname{Spec} E[t,t^{-1}] \cong T_E$ which we will use in the sequel to identify $X_*(T_E)$ with $\mathbb{Z}$. The inertia group $I = \mathrm{Gal}(E/K) = \{1,\gamma\}$ acts on $X_*(T) = \mathbb{Z}$ via $\gamma(\mu) = -\mu$ and hence $X_*(T)_I = \mathbb{Z}/2\mathbb{Z}$.

Consider the Néron-model $\mathcal{T} = \ker(N_{\mathcal{O}_E/\mathcal{O}_K}\colon \operatorname{Res}_{\mathcal{O}_E/\mathcal{O}_K}\mathbb{G}_m \to \mathbb{G}_m)$. As a scheme it is isomorphic to $\operatorname{Spec}\mathbb{F}_q[\![z]\!][a,b]/(a^2 - b^2 z - 1)$. Its special fiber has two connected components distinguished by $a \equiv 1$ or $-1 \bmod z$. Therefore the connected component of identity of $\mathcal{T}$ is $\mathcal{T}^0 := \operatorname{Spec}\mathbb{F}_q[\![z]\!][a',b]/(2a' + z(a')^2 - b^2)$, where $a = 1 + za'$. In particular $(-1,0) \notin \mathcal{T}^0\big(\mathbb{F}_q(\!(z)\!)\big)$. We let $\mathbb{P}$ be the group scheme $\mathcal{T}^0$ over $\mathbb{D} = \operatorname{Spec}\mathbb{F}_q[\![z]\!]$. Here $\mathbb{F} = \mathbb{F}_q$ and $\hat{\sigma} = \sigma$ is the q-Frobenius.

With each element of $X_*(T)_I$ the inverse of the Kottwitz map $\kappa_T\colon LT(\mathbb{F}_q^{\mathrm{alg}}) \to X_*(T)_I$, see [Kot85, 2.5], associates a $\hat{\sigma}$-conjugacy class in $LT(\mathbb{F}_q^{\mathrm{alg}})$. For example for $\bar{\mu} = 1 \in X_*(T)_I = \mathbb{Z}/2\mathbb{Z}$ one has to choose a lift $\mu \in X_*(T_E)$. If we choose $\mu = 1$ then with $\bar{\mu} = 1$ it associates

$$\begin{aligned} N_{E/K}(\mu(y)) &= \mu(y)\cdot\gamma(\mu(y)) \\ &= \Big(\frac{1}{2}(y + y^{-1}), \frac{1}{2y}(y^{-1} - y)\Big)\cdot\Big(\frac{1}{2}(-y - y^{-1}), \frac{1}{-2y}(-y^{-1} - (-y))\Big) \\ &= (-1,0) \in T(\mathbb{F}_q^{\mathrm{alg}}(\!(z)\!)). \end{aligned}$$

The $\hat{\sigma}$-conjugacy class given by $(-1,0)$ is independent of the choice of μ and of the uniformizer y (and of E) by [Kot85, 2.5].

Now consider the local $\mathcal{T}^0$-shtuka $\big((L^+\mathcal{T}^0)_{\mathbb{F}_q}, (-1,0)\big)$ over $\mathbb{F}_q$ which is bounded by $1 \in \widetilde{W} = X_*(T)_I$. We want to lift it to a local $\mathcal{T}^0$-shtuka over $\mathbb{F}_q[\![\zeta]\!][\xi]/(\xi^2 - \zeta)$ which is bounded. (It turns out that such a lift only exists over a ramified extension of $\mathbb{F}_q[\![\zeta]\!]$.)

Consider the isomorphism $K = \mathbb{F}_q(\!(z)\!) \to \mathbb{F}_q(\!(\zeta)\!), z \mapsto \zeta$. Fix an embedding $i\colon E \hookrightarrow \mathbb{F}_q(\!(\zeta)\!)^{\mathrm{alg}}$, set $\xi := i(y)$ and lift $N_{E/K}(\mu(y))$ to $g(E,\mu,i) := N_{E/K}\big(\mu\big(y - i(y)\big)\big)$. For example if $\mu = 1 \in \mathbb{Z} = X_*(T_E)$ we compute

$$(\alpha,\beta) \;:=\; g(E,1,i) \;=\; \mu(y-\xi)\cdot\gamma(\mu(y-\xi)) \;=$$

$$\left(\frac{(y-\xi) + (y-\xi)^{-1}}{2}, \frac{(y-\xi)^{-1} - (y-\xi)}{2y}\right)\cdot\gamma\left(\frac{(y-\xi) + (y-\xi)^{-1}}{2}, \frac{(y-\xi)^{-1} - (y-\xi)}{2y}\right),$$

with $\gamma(y) = -y$ and $\gamma|_{\mathbb{F}_q(\!(\xi)\!)} = \mathrm{id}$. Then

$$\begin{aligned}\alpha &= \tfrac{1}{4}\left((y-\xi)+(y-\xi)^{-1}\right)\left((-y-\xi)+(-y-\xi)^{-1}\right)\\ &\quad -\tfrac{1}{4}\left((y-\xi)^{-1}-(y-\xi)\right)\left((-y-\xi)^{-1}-(-y-\xi)\right)\\ &= \frac{1}{2}\left(\frac{(y-\xi)^2+(-y-\xi)^2}{(-y-\xi)(y-\xi)}\right)\\ &= \frac{\zeta+z}{\zeta-z},\end{aligned}$$

and

$$\begin{aligned}\beta &= \tfrac{-1}{4y}\left((y-\xi)+(y-\xi)^{-1}\right)\left((-y-\xi)^{-1}-(-y-\xi)\right)\\ &\quad +\tfrac{1}{4y}\left((y-\xi)^{-1}-(y-\xi)\right)\left((-y-\xi)+(-y-\xi)^{-1}\right)\\ &= \frac{2\xi}{\zeta-z}.\end{aligned}$$

Thus we get

$$g(E,1,i) = N_{E/K}(\mu(y-\xi)) = \left(\frac{\zeta+z}{\zeta-z}, \frac{2\xi}{\zeta-z}\right).$$

This shows that we can lift $\left((L^+\mathcal{T}^0)_{\mathbb{F}_q}, (-1,0)\right)$ to a local $\mathcal{T}^0$-shtuka $\left((L^+\mathcal{T}^0)_{\mathbb{F}_q[\![\xi]\!]}, g(E,\mu,i)\right)$ over $\mathbb{F}_q[\![\xi]\!]$ which is bounded by the bound $\hat{Z}_{\mu,i} := L^+\mathcal{T}^0 \cdot g(E,\mu,i) \cdot L^+\mathcal{T}^0/L^+\mathcal{T}^0 \subset \mathcal{F}\ell_{\mathcal{T}^0} \widehat{\times}_{\operatorname{Spec}\mathbb{F}_q} \operatorname{Spf}\mathbb{F}_q[\![\xi]\!]$. However the lift, and as we will see also the bound, depends on the choice of μ and of the embedding i.

We first compute how $g(E,\mu,i)$ depends on the chosen embedding i. For this purpose we compute $\dfrac{N_{E/K}(\mu(y-i(y)))}{N_{E/K}(\mu(y-i\circ\gamma(y))}$. Changing i to $i\circ\gamma$ replaces ξ by $-\xi$ and we have

$$g(E,1,i\circ\gamma) = N_{E/K}(\mu(y+\xi)) = \left(\frac{\zeta+z}{\zeta-z}, \frac{-2\xi}{\zeta-z}\right).$$

Note that $g(E,1,i) = g(E,1,i\circ\gamma)^{-1}$. Therefore

$$\frac{g(E,1,i)}{g(E,1,i\circ\gamma)} = g(E,1,i)^2 = \left(\frac{(z+\zeta)^2+4\zeta z}{(\zeta-z)^2}, \frac{4\xi(z+\zeta)}{(\zeta-z)^2}\right).$$

This also shows what happens if we replace $\mu \in X_*(T_E) = \mathbb{Z}$ by another lift of $\bar{\mu} \in \mathbb{Z}/2\mathbb{Z}$, i.e.

$$\frac{g(E,\mu+2,i)}{g(E,\mu,i)} = g(E,2,i) = g(E,1,i)^2.$$

Observe that $\dfrac{g(E,1,i)}{g(E,1,i\circ\gamma)} \in \mathcal{T}^0(\mathbb{F}_q[\![z,\tfrac{\xi}{z}]\!]) \smallsetminus \mathcal{T}^0(\mathbb{F}_q[\![\xi,z]\!])$. So $g(E,1,\gamma\circ i)$ does not lie in the closure of the subscheme $L^+\mathcal{T}^0 \cdot g(E,1,i) \cdot L^+\mathcal{T}^0 \subset L\mathcal{T}^0 \widehat{\times}_{\operatorname{Spec}\mathbb{F}_q} \operatorname{Spf}\mathbb{F}_q[\![\xi]\!]$. In particular the bounds defined by $\hat{Z}_{\mu,i} := L^+\mathcal{T}^0 \cdot g(E,\mu,i) \cdot L^+\mathcal{T}^0/L^+\mathcal{T}^0 \subset \mathcal{F}\ell_{\mathcal{T}^0} \widehat{\times}_{\operatorname{Spec}\mathbb{F}_q} \operatorname{Spf}\mathbb{F}_q[\![\xi]\!]$ depend on the embedding $i\colon E \hookrightarrow \mathbb{F}_q(\!(\zeta)\!)^{\mathrm{alg}}$ and the lift $\mu \in X_*(T)$ of $\bar{\mu} \in X_*(T)_I$. For this reason we decided to treat bounds axiomatically in Definition 4.5.

4.3 Representability of the Bounded Rapoport-Zink Functor

Let P denote the generic fiber of $\mathbb{P}$ and let $b \in LP(k)$ for some field $k \in \mathcal{N}ilp_{\mathbb{F}[\![\zeta]\!]}$. With b Kottwitz associates a slope homomorphism

$$\nu_b\colon D_{k(\!(z)\!)} \to P_{k(\!(z)\!)},$$

called Newton polygon of b; see [Kot85, 4.2]. Here D is the diagonalizable pro-algebraic group over $k(\!(z)\!)$ with character group $\mathbb{Q}$. The slope homomorphism is characterized by assigning the slope filtration of $(V \otimes_{\mathbb{F}(\!(z)\!)} k(\!(z)\!), \rho(b) \cdot (\operatorname{id} \otimes \hat{\sigma}))$ to any $\mathbb{F}(\!(z)\!)$-rational representation (V, ρ) of P; see [Kot85, Section 3]. We assume that $b \in LP(k)$ satisfies a *decency equation for a positive integer* s, that is,

$$(b\hat{\sigma})^s \;=\; s\nu_b(z)\,\hat{\sigma}^s \qquad \text{in} \quad LP(k) \ltimes \langle \hat{\sigma} \rangle. \tag{4.3}$$

Remark 4.10. Assume that $b \in LP(k)$ is decent with the integer s and let $\ell \subset k^{\text{alg}}$ be the finite field extension of $\mathbb{F}$ of degree s. Then $b \in LP(\ell)$ because by (4.3) the element b has values in the fixed field of $\hat{\sigma}^s$ which is ℓ. Note that if k is algebraically closed, any $\hat{\sigma}$-conjugacy class in $LP(k)$ contains an element satisfying a decency equation; see [Kot85, Section 4].

Remark 4.11. With the element $b \in LP(k)$ one can associate a connected algebraic group J_b over $\mathbb{F}(\!(z)\!)$ which is defined by its functor of points that assigns to an $\mathbb{F}(\!(z)\!)$-algebra R the group

$$J_b(R) := \big\{ g \in P(R \otimes_{\mathbb{F}(\!(z)\!)} k(\!(z)\!)) \colon g^{-1} b \hat{\sigma}(g) = b \big\}.$$

Let b satisfy a decency equation for the integer s and let F_s be the fixed field of $\hat{\sigma}^s$ in $k(\!(z)\!)$. Then ν_b is defined over F_s and $J_b \times_{\mathbb{F}(\!(z)\!)} F_s$ is the centralizer of the 1-parameter subgroup $s\nu_b$ of P and hence a Levi subgroup of P_{F_s}; see [RZ96, Corollary 1.9]. In particular $J_b(\mathbb{F}(\!(z)\!)) \subset P(F_s) \subset LP(\ell)$ where ℓ is the finite field extension of $\mathbb{F}$ of degree s.

Let $\hat{Z} \subset \widehat{\mathcal{F}\ell}_{\mathbb{P}}$ be a bound (see Definition 4.5) and set $Z = \hat{Z} \hat{\times}_{\operatorname{Spf} \mathbb{F}[\![\zeta]\!]} \operatorname{Spec} \mathbb{F}$. We define the associated *affine Deligne-Lusztig variety* as the reduced closed ind-subscheme $X_Z(b) \subset \mathcal{F}\ell_{\mathbb{P}}$ whose K-valued points (for any field extension K of k) are given by

$$X_Z(b)(K) := \big\{ g \in \mathcal{F}\ell_{\mathbb{P}}(K) \colon g^{-1}\, b\, \hat{\sigma}^*(g) \in Z(K) \big\}.$$

If $\omega \in \widetilde{W}$ and $Z = \mathcal{S}(\omega)$ is the Schubert variety from Example 4.8, we set $X_\omega(b) := X_{\mathcal{S}(\omega)}(b)$.

Definition 4.12. Let $\underline{\mathcal{L}}_0$ be a local $\mathbb{P}$-shtuka over an $\mathbb{F}$-scheme $\bar{T}$. Consider the functor $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}$ from Definition 4.1 and its subfunctor

$$\begin{aligned} \underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}^{\hat{Z}} \colon (\mathcal{N}ilp_{k[\![\zeta]\!]})^o \quad &\longrightarrow \quad \mathcal{S}ets \\ S \quad &\longmapsto \quad \Big\{ \text{Isomorphism classes of } (\underline{\mathcal{L}}, \bar{\delta}) \in \underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}(S) \colon \underline{\mathcal{L}} \text{ is bounded by } \hat{Z} \Big\} \end{aligned}$$

$\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}$ is an ind-scheme over $\operatorname{Spf} k[\![\zeta]\!]$ by Theorem 4.4, and $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}^{\hat{Z}}$ is a closed ind-subscheme by Proposition 4.7.

In the remaining part of the chapter we prove that $\underline{\mathcal{M}}_{\underline{\mathcal{L}}_0}^{\hat{Z}}$ is pro-representable by a formal scheme in the following important special situation. Let $\bar{T} = \operatorname{Spec} k$ and let $\underline{\mathbb{L}}_0 = (L^+\mathbb{P}, b\hat{\sigma}^*)$ be a trivialized local $\mathbb{P}$-shtuka over $\operatorname{Spec} k$. Assume that b is decent with integer s and let $\ell \subset k^{\text{alg}}$ be the finite field extension of $\mathbb{F}$ of degree s. Then $b \in LP(\ell)$ by Remark 4.10. So $\underline{\mathbb{L}}_0$ is defined over $\bar{T} = \operatorname{Spec} \ell$ and we may replace k by ℓ. Then $\underline{\mathcal{M}}_{\underline{\mathbb{L}}_0}^{\hat{Z}}$ is defined over $\operatorname{Spf} \ell[\![\zeta]\!]$.

Theorem 4.13. *The functor* $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}\colon (\mathcal{N}ilp_{\ell[\![\zeta]\!]})^o \to \mathcal{S}ets$ *is pro-representable by a formal scheme over* $\operatorname{Spf}\ell[\![\zeta]\!]$ *which is locally formally of finite type. Its underlying reduced subscheme equals* $X_Z(b)$*. In particular* $X_Z(b)$ *is a scheme locally of finite type over* ℓ*. The formal scheme representing* $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}$ *is called a* bounded Rapoport-Zink space for local $\mathbb{P}$-shtukas*.*

Recall that a formal scheme over $\ell[\![\zeta]\!]$ in the sense of [EGA, I_{new}, 10] is called *locally formally of finite type* if it is locally noetherian and adic and its reduced subscheme is locally of finite type over ℓ. It is called *formally of finite type* if in addition it is quasi-compact.

Remark 4.14. By our assumptions $\mathrm{QIsog}_\ell(\underline{\mathbb{L}}_0)$ equals the group $J_b(\mathbb{F}(\!(z)\!))$ from Remark 4.11. This group acts on the functor $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}$ via $g\colon (\underline{\mathcal{L}}, \bar\delta) \mapsto (\underline{\mathcal{L}}, g\circ\bar\delta)$ for $g \in \mathrm{QIsog}_\ell(\underline{\mathbb{L}}_0)$.

Proof of Theorem 4.13. The proof will use a sequence of lemmas and will eventually be complete after Lemma 4.21. Consider the universal local $\mathbb{P}$-shtuka $\underline{\mathcal{L}}_{univ}$ over $\widehat{\mathcal{F}\ell}_{\mathbb{P},\ell} := \mathcal{F}\ell_{\mathbb{P}} \widehat{\times}_{\mathbb{F}} \operatorname{Spf}\ell[\![\zeta]\!]$ (see Theorem 4.4). Let $\mathcal{M}_{\mathbb{L}_0}^{\hat{Z}}$ be the closed ind-subscheme of $\widehat{\mathcal{F}\ell}_{\mathbb{P},\ell}$ over which $\underline{\mathcal{L}}_{univ}$ is bounded by $\hat{Z}$. By construction $\mathcal{M}_{\mathbb{L}_0}^{\hat{Z}}$ pro-represents the functor $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}$. It is clear that the reduced ind-subscheme equals $X_Z(b)$. We have to show that $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}$ is locally formally of finite type. By rigidity of quasi-isogenies the functor $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}$ is equivalent to the following functor

$$\begin{aligned} (\mathcal{N}ilp_{\ell[\![\zeta]\!]})^o \quad &\to \quad \mathcal{S}ets \\ S \quad &\longmapsto \quad \Big\{ (\underline{\mathcal{L}}, \delta)\colon \underline{\mathcal{L}} \text{ is a local } \mathbb{P}\text{-shtuka over } S \text{ bounded by } \hat{Z} \\ &\qquad\qquad \text{and } \delta\colon \underline{\mathcal{L}} \to \underline{\mathbb{L}}_{0,S} \text{ is a quasi-isogeny} \Big\}/\sim . \end{aligned}$$

We take a representation $\iota\colon \mathbb{P} \to \mathrm{SL}_{r,\mathbb{D}} =: H$ with quasi-affine quotient $H/\mathbb{P}$; see Proposition 2.13(a). It induces a 1-morphism

$$\iota_* := \mathscr{H}^1(\iota)\colon \mathscr{H}^1(\operatorname{Spec}\mathbb{F}, L^+\mathbb{P})(S) \to \mathscr{H}^1(\operatorname{Spec}\mathbb{F}, L^+H)(S).$$

For an $L^+\mathbb{P}$-torsor $\mathcal{L}_+$ over S we denote by $\mathcal{V}(\iota_*\mathcal{L}_+)$ the sheaf of $\mathcal{O}_S[\![z]\!]$-modules associated with the image of $\mathcal{L}$ in $\mathscr{H}^1(\operatorname{Spec}\mathbb{F}, L^+\,\mathrm{GL}_r)(S)$ by Remark 3.2. In particular $\mathcal{V}(\iota_*(L^+\mathbb{P})_S) = \mathcal{O}_S[\![z]\!]^{\oplus r}$. Let $\rho^\vee$ be the half-sum of all positive coroots of H.

For each non-negative integer $n \in \mathbb{N}_0$ let $\mathcal{M}^n$ be the closed ind-subscheme of $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}$ defined by the following sub functor of $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}$

$$\begin{aligned} \underline{\mathcal{M}}^n\colon (\mathcal{N}ilp_{\ell[\![\zeta]\!]})^o \quad &\longrightarrow \quad \mathcal{S}ets \\ S \quad &\longmapsto \quad \Big\{ (\underline{\mathcal{L}}, \delta) \in \underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}(S)\colon \mathscr{H}^1(\iota)(\delta) \text{ is bounded by } 2n\rho^\vee \Big\}, \end{aligned}$$

where $\underline{\mathcal{L}} = (\mathcal{L}_+, \hat\tau)$ and where we say that $\mathscr{H}^1(\iota)(\delta)$ *is bounded by* $2n\rho^\vee$ if for all $j = 1, \ldots, r$

$$\textstyle\bigwedge^j_{\mathcal{O}_S[\![z]\!]} \mathscr{H}^1(\iota)(\delta)\big(\mathcal{V}(\iota_*\mathcal{L}_+)\big) \;\subset\; z^{n(j^2-jr)} \cdot \textstyle\bigwedge^j_{\mathcal{O}_S[\![z]\!]} \mathcal{V}(\iota_*(L^+\mathbb{P})_S)\,. \tag{4.4}$$

By [HV11, Lemma 4.3] the latter is equivalent to the boundedness condition considered in [HV11, Definition 3.5], see Example 4.9, because $\rho^\vee = (r-1, \ldots, 1-r)$ and (4.4) is automatically an equality for $j = r$ as ι factors through H.

Lemma 4.15. *The ind-scheme* $\mathcal{M}^n$ *representing the functor* $\underline{\mathcal{M}}^n$ *is a* ζ*-adic noetherian formal scheme over* $\ell[\![\zeta]\!]$ *whose underlying topological space* $(\mathcal{M}^n)_{\mathrm{red}}$ *is a quasi-projective scheme over* $\operatorname{Spec}\ell$*.*

Proof. Since $H/\mathbb{P}$ is quasi-affine, the induced morphism $\iota_*\colon \widehat{\mathcal{F}\ell}_{\mathbb{P},\ell} \to \widehat{\mathcal{F}\ell}_{H,\ell}$ is a locally closed embedding by Remark 4.3. The representation ι induces a functor ι_* from local $\mathbb{P}$-shtukas to local H-shtukas. Consider the local H-shtuka $\underline{\mathbb{H}}_0 := \iota_*\underline{\mathbb{L}}_0 = ((L^+H)_\ell, \iota(b)\hat{\sigma}^*)$ over ℓ and view $\widehat{\mathcal{F}\ell}_{H,\ell}$ as a moduli space representing the functor $\underline{\mathcal{M}}_{\underline{\mathbb{H}}_0}$, parametrizing local H-shtukas together with a quasi-isogeny δ_H to $\underline{\mathbb{H}}_0$; see Theorem 4.4. Let $\widehat{\mathcal{F}\ell}^{(n)}_{H,\ell}$ be the closed ind-subscheme of $\widehat{\mathcal{F}\ell}_{H,\ell}$ defined by condition (4.4), that is, by bounding δ_H by $2n\rho^\vee$. It is a ζ-adic noetherian formal scheme over $\mathrm{Spf}\,\ell[\![\zeta]\!]$ by [HV11, Proposition 5.5] whose underlying topological space is a projective scheme over $\mathrm{Spec}\,\ell$. Therefore

$$\mathcal{M}^n(i) \;:=\; \mathcal{M}^n \,\widehat{\times}_{\mathrm{Spf}\,\ell[\![\zeta]\!]}\, \mathrm{Spec}\,\ell[\![\zeta]\!]/(\zeta^i) \;=\; \underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0} \,\widehat{\times}_{\widehat{\mathcal{F}\ell}_{H,\ell}}\, \widehat{\mathcal{F}\ell}^{(n)}_{H,\ell} \,\widehat{\times}_{\mathrm{Spf}\,\ell[\![\zeta]\!]}\, \mathrm{Spec}\,\ell[\![\zeta]\!]/(\zeta^i)$$

is a scheme of finite type over $\mathrm{Spec}\,\ell[\![\zeta]\!]/(\zeta^i)$ for all i with underlying topological space $\mathcal{M}^n(1)$ independent of i. Now our claim follows from [EGA, $\mathrm{I}_{\mathrm{new}}$, Cor 10.6.4]. ☐

For each non-negative integer $n \in \mathbb{N}_0$ we define the following sub functor of $\underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0}$

$$\begin{array}{rcl} \underline{\mathcal{M}}_n\colon (\mathcal{N}ilp_{\ell[\![\zeta]\!]})^o & \longrightarrow & \mathcal{S}ets \\ S & \longmapsto & \Big\{ (\underline{\mathcal{L}}, \delta) \in \underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0}(S)\colon \text{ for any closed point } s \text{ in } S, \\ & & \qquad \mathscr{H}(\iota)(\delta_s) \text{ is bounded by } 2n\rho^\vee \Big\}. \end{array}$$

This functor is represented by an ind-scheme $\mathcal{M}_n$ which is the formal completion of $\underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0}$ along the quasi-compact scheme $(\mathcal{M}^n)_{red}$.

Lemma 4.16. *$\mathcal{M}_n$ is a formal scheme formally of finite type over* $\mathrm{Spf}\,\ell[\![\zeta]\!]$.

To prove the lemma we need to start with the following definition.

Definition 4.17. Let $R = \varprojlim R_m \in \mathcal{N}ilp_{\mathbb{F}[\![\zeta]\!]}$ for a projective system $(R_m, u_{m,m'})$ of discrete rings indexed by $\mathbb{N}_0$. Suppose that all homomorphisms $R \to R_m$ are surjective, and that all the kernels $I_m := \ker(u_{m,0}\colon R_m \to R_0) \subset R_m$ are nilpotent. A local $\mathbb{P}$-shtuka over $\mathrm{Spf}\,R$ is a projective system $(\underline{\mathcal{L}}_m)_{m\in\mathbb{N}_0}$ of local $\mathbb{P}$-shtukas $\underline{\mathcal{L}}_m$ over R_m with $\underline{\mathcal{L}}_{m-1} \cong \underline{\mathcal{L}}_m \otimes_{R_m} R_{m-1}$.

Lemma 4.18. *Let R in $\mathcal{N}ilp_{\mathbb{F}[\![\zeta]\!]}$ be as in the above definition. The pull back functor defines an equivalence between the category of local $\mathbb{P}$-shtukas over* $\mathrm{Spec}\,R$ *bounded by $\hat{Z}$ and the category of local $\mathbb{P}$-shtukas over* $\mathrm{Spf}\,R$ *bounded by $\hat{Z}$.*

Proof. Since R is in $\mathcal{N}ilp_{\mathbb{F}[\![\zeta]\!]}$ there is an integer $e \in \mathbb{N}$ such that $\zeta^e = 0$ on R. Let $\underline{\mathcal{L}} := (\underline{\mathcal{L}}_m)_{m\in\mathbb{N}_0}$ be a local $\mathbb{P}$-shtuka over $\mathrm{Spf}\,R$. There is an étale covering $R'_0 \to R_0$ which trivializes $\underline{\mathcal{L}}_0$. By [SGA 1, Théorème I.8.3] there is a unique étale R-algebra R' with $R' \otimes_R R_0 \cong R'_0$. As in [HV11, Proposition 2.2(c)] this gives rise to compatible trivializations $\underline{\mathcal{L}}_m \otimes_{R_m} R'_m \cong ((L^+\mathbb{P})_{R'_m}, b_m\hat{\sigma}^*)$ over $R'_m := R' \otimes_R R_m$ for all m. Here $b_m \in LP(R'_m)$ and $b_m \otimes_{R'_m} R'_{m-1} = b_{m-1}$.

Take a faithful representation $\mathbb{P} \hookrightarrow \mathrm{SL}_{r,\mathbb{D}}$ and consider the induced closed immersion $LP \hookrightarrow L\,\mathrm{SL}_{r,\mathbb{D}}$. The ind-scheme structure on $\mathcal{F}\ell_{\mathrm{SL}_{r,\mathbb{D}}}$ is given by $\mathcal{F}\ell_{\mathrm{SL}_{r,\mathbb{D}}} = \varinjlim \mathcal{F}\ell^{(n)}_{\mathrm{SL}_{r,\mathbb{D}}}$ where $\mathcal{F}\ell^{(n)}_{\mathrm{SL}_{r,\mathbb{D}}}$ is defined by condition (4.4). Let $L\,\mathrm{SL}^{(n)}_{r,\mathbb{D}} = L\,\mathrm{SL}_{r,\mathbb{D}} \times_{\mathcal{F}\ell_{\mathrm{SL}_{r,\mathbb{D}}}} \mathcal{F}\ell^{(n)}_{\mathrm{SL}_{r,\mathbb{D}}}$. Then

$$L\,\mathrm{SL}^{(n)}_{r,\mathbb{D}}(S) \;=\; \big\{\, g \in L\,\mathrm{SL}_{r,\mathbb{D}}(S)\colon \text{ all } j \times j\text{-minors of } g \text{ lie in } z^{-n(j^2-jr)}\mathcal{O}_S(S)[\![z]\!] \;\;\forall\, j \,\big\}.$$

This implies that $L\,\mathrm{SL}^{(n)}_{r,\mathbb{D}}$, and hence also $LP^{(n)} := LP \times_{L\,\mathrm{SL}_{r,\mathbb{D}}} L\,\mathrm{SL}^{(n)}_{r,\mathbb{D}}$ is an infinite dimensional affine scheme over $\mathbb{F}$. Since $Z_e := \hat{Z}\,\widehat{\times}_{\mathrm{Spf}\,\mathbb{F}[\![\zeta]\!]}\,\mathrm{Spec}\,\mathbb{F}[\![\zeta]\!]/(\zeta^e)$ has the same underlying topological space as Z, it is quasi-compact. So there is an $n \in \mathbb{N}$ such that $Z_e \subseteq \mathcal{F}\ell^{(n)}_{\mathrm{SL}_{r,\mathbb{D}}} \times_{\mathrm{Spec}\,\mathbb{F}} \mathrm{Spec}\,\mathbb{F}[\![\zeta]\!]/(\zeta^e)$. As one sees from the following diagram

$$\begin{tikzcd}
\mathrm{Spec}\,R'_m \arrow[r, dashed, "b_m"] \arrow[d] & L\,\mathrm{SL}^{(n)}_{r,\mathbb{D}} \times_{\mathbb{F}} \mathrm{Spec}\,\mathbb{F}[\![\zeta]\!]/(\zeta^e) \arrow[r] \arrow[d] & L\,\mathrm{SL}_{r,\mathbb{D}} \times_{\mathbb{F}} \mathrm{Spec}\,\mathbb{F}[\![\zeta]\!]/(\zeta^e) \arrow[d] \\
Z_e \arrow[r] & \mathcal{F}\ell^{(n)}_{\mathrm{SL}_{r,\mathbb{D}}} \times_{\mathbb{F}} \mathrm{Spec}\,\mathbb{F}[\![\zeta]\!]/(\zeta^e) \arrow[r] & \mathcal{F}\ell_{\mathrm{SL}_{r,\mathbb{D}}} \times_{\mathbb{F}} \mathrm{Spec}\,\mathbb{F}[\![\zeta]\!]/(\zeta^e)\,,
\end{tikzcd}$$

the morphism $b_m\colon \mathrm{Spec}\,R'_m \to LP$ factors through $LP^{(n)} = LP \times_{L\,\mathrm{SL}_{r,\mathbb{D}}} L\,\mathrm{SL}^{(n)}_{r,\mathbb{D}}$ for all m. Since $LP^{(n)}$ is affine, the compatible collection of morphisms $b_m\colon \mathrm{Spec}\,R'_m \to LP^{(n)}$ is given by a compatible collection of homomorphisms $\mathcal{O}(LP^{(n)}) \to R'_m$. It corresponds to a homomorphism $\mathcal{O}(LP^{(n)}) \to R'$, because $R' = \varprojlim R'_m$, and thus to a morphism $b_\infty\colon \mathrm{Spec}\,R' \to LP^{(n)}$. This gives the local $\mathbb{P}$-shtuka $((L^+\mathbb{P})_{R'}, b_\infty\hat{\sigma}^*)$ over $\mathrm{Spec}\,R'$ which carries a descent datum from the $\underline{\mathcal{L}}_m$ and hence induces a local $\mathbb{P}$-shtuka over $\mathrm{Spec}\,R$. ☐

Let us come back to the

Proof of Lemma 4.16. For each $m \geq n$ let $\mathcal{M}^m_n$ be the formal completion of $\mathcal{M}^m$ along $(\mathcal{M}_n)_{\mathrm{red}}$. It is an adic formal scheme over $\ell[\![\zeta]\!]$. Let U be an affine open subscheme of $(\mathcal{M}_n)_{\mathrm{red}}$. This defines an affine open formal subscheme $\mathrm{Spf}\,R_m$ of $\mathcal{M}^m_n$ with underlying set U. Let R be the inverse limit of the projective system $R_{m+1} \to R_m$ and let $\mathfrak{a}_m$ denote the ideal such that $R_m = R/\mathfrak{a}_m$. Let J be the inverse image in R of the largest ideal of definition in R_n. We want to show that R is J-adic. We make the following

Claim. For any integer $c > 0$ there is an integer m_0 such that for any $m \geq m_0$ the natural map $R_m/J^cR_m \to R_{m_0}/J^cR_{m_0}$ is an isomorphism.

To prove the claim let $\underline{\mathcal{L}}_m$ be the universal local $\mathbb{P}$-shtuka over $\mathrm{Spf}\,R_m$. Consider the local $\mathbb{P}$-shtuka $(\underline{\mathcal{L}}_m)_m$ over $\mathrm{Spf}\,R$ and its pullback over $\mathrm{Spf}\,R/J^c$. The latter comes from a local $\mathbb{P}$-shtuka $\underline{\mathcal{L}}$ over $\mathrm{Spec}\,R/J^c \in \mathcal{N}ilp_{\ell[\![\zeta]\!]}$ by Lemma 4.18. By rigidity of quasi-isogenies the quasi-isogeny δ_n over $R/J = R_n/J$ lifts to a quasi-isogeny δ over R/J^c. Since $\mathrm{Spec}\,R/J^c$ is quasi-compact, the quasi-isogeny $\mathscr{H}^1(\iota)(\delta)$ satisfies condition (4.4) for some m_0, that is, it is bounded by $2m_0\rho^\vee$. By the universal property of $\mathcal{M}^{m_0}_n$ the tuple $(\underline{\mathcal{L}}, \delta)$ induces a morphism $R_{m_0} \to R/J^c$ making the following diagram commutative, from which the claim follows

$$\begin{tikzcd}
R \arrow[r, two heads] \arrow[d] & R_m \arrow[r, two heads] \arrow[d] & R_{m_0} \arrow[d] \arrow[dll, dashed] \\
R/J^c \arrow[r, two heads] & R_m/J^cR_m \arrow[r, two heads] & R_{m_0}/J^cR_{m_0}.
\end{tikzcd}$$

The claim implies that the chain $\mathfrak{a}_n + J^c \supseteq \mathfrak{a}_{n+1} + J^c \supseteq \ldots \supseteq \mathfrak{a}_i + J^c \supseteq \ldots$ stabilizes. Now set $\mathcal{J}_c := \bigcap_i \mathfrak{a}_i + J^c = \mathfrak{a}_m + J^c$ for $m \gg 0$ and consider the descending chain $\mathcal{J}_1 \supseteq \mathcal{J}_2 \supseteq \ldots$. Note that $\mathcal{J}_1 = J$ and $\mathcal{J}_{c+1} + \mathcal{J}_1^c = \mathcal{J}_c$. Since $\mathcal{J}_1/\mathcal{J}_2 \cong JR_m/J^2R_m$ for $m \gg 0$, it is a finitely generated R-module. Therefore $\mathcal{M}_n$ is a locally noetherian adic formal scheme over $\mathrm{Spf}\,\ell[\![\zeta]\!]$ by [RZ96, Proposition 2.5]. It is formally of finite type because $(\mathcal{M}_n)_{\mathrm{red}} = (\mathcal{M}^n)_{\mathrm{red}}$ is quasi-projective over ℓ by Lemma 4.15. ☐

Remark 4.19. Up to this point we have not used that $\underline{\mathbb{L}}_0$ is decent and, in fact, Lemmas 4.15 and 4.16 also hold for the ind-scheme $\underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathcal{L}}_0}$ in the more general situation where $\underline{\mathcal{L}}_0$ is a (not necessarily trivialized) local $\mathbb{P}$-shtuka over an arbitrary $\mathbb{F}$-scheme $\bar{T}$.

From now on we use that $\bar{T} = \operatorname{Spec} \ell$ and that $\underline{\mathbb{L}}_0$ is decent with the integer s. In the sequel we consider points $x \in \underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0}(K)$ for varying field extensions K of ℓ. For two points $x := (\underline{\mathcal{L}}, \delta)$ and $x' := (\underline{\mathcal{L}}', \delta')$ in $\underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0}(K)$ we define

$$\tilde{d}(x, x') := \min\big\{n \in \mathbb{N}_0 \colon \mathscr{H}^1(\iota)(\delta^{-1}\delta') \text{ is bounded by } 2n\rho^\vee\big\}. \tag{4.5}$$

In the situation where $\delta = g$ and $\delta' = g'$ for $g, g' \in LP(K)$, as well as $x = ((L^+\mathbb{P})_K, g^{-1}b\hat{\sigma}^*(g), g)$ and $x' = ((L^+\mathbb{P})_K, (g')^{-1}b\hat{\sigma}^*(g'), g')$, we also write $\tilde{d}(g, g') := \tilde{d}(x, x')$. Note that a point $x \in \underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0}(K)$ belongs to $\mathcal{M}_n$ if and only if $\tilde{d}((\underline{\mathbb{L}}_0, \mathrm{id}), x) \leq n$.

Although we will not use this, note that $\tilde{d}$ is a metric on $\underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0}$. This follows from the fact that (4.4) for $n = 0$ together with Cramer's rule implies that $\mathscr{H}^1(\iota)(\delta_1^{-1}\delta_2)$ is an isomorphism $\mathcal{V}(\iota_*\underline{\mathcal{L}}_2) \xrightarrow{\sim} \mathcal{V}(\iota_*\underline{\mathcal{L}}_1)$; compare the discussion around [AH13, Equations (3.3) and (3.4)].

Lemma 4.20. *For every integer $c > 0$ there is an integer $d_0 > 0$ with the following property. For every $g \in LP(K)$ with $\tilde{d}(g, b\,\hat{\sigma}^*(g)) < c$ there is a $g' \in LP(\ell)$ with $\tilde{d}(g, g') \leq d_0$.*

Proof. This is just a reformulation of [RZ99, Theorem 1.4 and Subsection 2.1] taking into account that by functorial properties of Bruhat-Tits buildings (see [Lan00]), ι induces an injective isometric map of Bruhat-Tits buildings $\mathscr{B}(P) \to \mathscr{B}(H)$. □

Lemma 4.21. *There is an integer $d_0 \in \mathbb{N}_0$ such that $max\big\{\tilde{d}(x, \underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0}(\ell)) \colon x \in \underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0}\big\} \leq d_0$.*

Proof. Let $x := (\underline{\mathcal{L}}^+, \delta)$ be a point in $\underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0}(K)$. Take a trivialization $(\underline{\mathcal{L}}^+, \delta) \cong ((L^+\mathbb{P})_K, h\hat{\sigma}^*, g)$ with $\delta = g \in LP(K)$ and $h = g^{-1}b\hat{\sigma}^*(g)$. Since the local $\mathbb{P}$-shtuka $((L^+\mathbb{P})_K, h\hat{\sigma}^*)$ is bounded by $\hat{Z}$ and Z is quasi-compact by definition of the boundedness condition (see Definition 4.5), we have $\tilde{d}(g, b\,\hat{\sigma}^*(g)) = \min\{n \in \mathbb{N}_0 \colon \mathscr{H}^1(\iota)(h) \text{ is bounded by } 2n\rho^\vee\} < c$, for some natural number c which is independent of x. Let $d_0 = d_0(c)$ be the integer from Lemma 4.20. Then there is a $g' \in LP(\ell)$ with $\tilde{d}(g, g') \leq d_0$. Now we consider the associated point $x' := ((L^+\mathbb{P})_\ell, g'^{-1}b\hat{\sigma}^*g', g')$ of $\underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0}(\ell)$. We conclude $\tilde{d}(x, x') \leq d_0$ and the lemma follows. □

For a point $y \in \underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0}(\ell)$ set $B(y, d_0) := \big\{x \in \underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0} : \tilde{d}(x, y) \leq d_0\big\}$. Here $x \in \underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0}(K)$ for a field extension K of ℓ. We set $B_n(y, d_0) = B(y, d_0) \cap \mathcal{M}_n$. Note that these are closed subsets.
For each integer r let

$$\mathscr{Z}_n^r = \bigcup_{y \in \underline{\mathcal{M}}^{\hat{Z}}_{\underline{\mathbb{L}}_0}(\ell),\, \tilde{d}((\underline{\mathbb{L}}_0, \mathrm{id}), y) \geq r} B_n(y, d_0)$$

If $y \notin \mathcal{M}_{n+d_0}$, that is $\tilde{d}((\underline{\mathbb{L}}_0, \mathrm{id}), y) > n + d_0$, and $x \in \mathcal{M}_n$, that is $\tilde{d}((\underline{\mathbb{L}}_0, \mathrm{id}), x) \leq n$, then

$$\tilde{d}(x, y) \geq \tilde{d}((\underline{\mathbb{L}}_0, \mathrm{id}), x) - n + \tilde{d}(x, y) \geq \tilde{d}((\underline{\mathbb{L}}_0, \mathrm{id}), y) - n > d_0$$

and thus $B_n(y, d_0) = \emptyset$. We get

$$\mathscr{Z}_n^r = \bigcup_{y \in \mathcal{M}_{n+d_0}(\ell), \tilde{d}((\underline{\mathbb{L}}_0, \mathrm{id}), y) \geq r} B_n(y, d_0).$$

Since $\big(\mathcal{M}_{n+d_0}\big)_{\mathrm{red}} = \big(\mathcal{M}^{n+d_0}\big)_{\mathrm{red}}$ is quasi-projective over ℓ by Lemma 4.15, this union is finite.

Let $\mathcal{U}_n^r$ be the open formal sub-scheme of $\mathcal{M}_n$ whose underlying reduced set is $\mathcal{M}_n \smallsetminus \mathcal{Z}_n^r$. We claim that the chain $\mathcal{U}_n^r \hookrightarrow \mathcal{U}_{n+1}^r \hookrightarrow \ldots$ of open formal sub-schemes of $\mathcal{M}_n$ stabilizes. By the definition of $\mathcal{M}_n$ it is enough to verify this on the underlying set of points. Suppose that there is some element $x \in \mathcal{U}_{n+1}^r \smallsetminus \mathcal{U}_n^r$. By Lemma 4.21 there exist a $y \in \underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}(\ell)$ such that $\tilde{d}(x,y) \le d_0$. Since $x \notin \mathcal{Z}_{n+1}^r$ we must have $\tilde{d}((\underline{\mathbb{L}}_0, \mathrm{id}), y) < r$. Then

$$\tilde{d}((\underline{\mathbb{L}}_0, \mathrm{id}), x) \le \tilde{d}((\underline{\mathbb{L}}_0, \mathrm{id}), y) + \tilde{d}(x,y) < r + d_0, \tag{4.6}$$

therefore if $n \ge r + d_0$ then $\tilde{d}((\underline{\mathbb{L}}_0, \mathrm{id}), x) \le n$ which is a contradiction and consequently there is no such x.

Let $\mathcal{U}^r = \bigcup_n \mathcal{U}_n^r$ (which equals $\mathcal{U}_n^r$ for $n \ge r + d_0$). Note that every point x of $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}$ lies in the union of the $\mathcal{U}^r$s. Indeed, if $\tilde{d}\big((\underline{\mathbb{L}}_0, \mathrm{id}), x\big) < r - d_0$ for some r, then x is contained in $\mathcal{U}^r$, because otherwise there is a $y \in \underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}(\ell)$ with $\tilde{d}(x,y) \le d_0$ and $\tilde{d}\big((\underline{\mathbb{L}}_0, \mathrm{id}), y\big) \ge r$, a contradiction. Now consider the chain

$$\mathcal{U}^r \hookrightarrow \mathcal{U}^{r+1} \ldots \hookrightarrow \underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}$$

of open immersions of formal schemes formally of finite type, note that $\mathcal{U}^r$ is open in $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}$. Indeed the underlying topological space of $\mathcal{U}^r$ is open in $\mathcal{M}_n$ for every n and the ind-scheme $\mathcal{M}_{\mathbb{L}_0}^{\hat{Z}}$ carries the limit topology of the limit over the $\mathcal{M}_n$. This shows that the formal scheme $\mathcal{U}^r$ equals the formal completion of the open ind-scheme $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}|_{|\mathcal{U}^r|}$ of $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}$ supported on $|\mathcal{U}^r|$ along the whole set $|\mathcal{U}^r|$ and thus $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}|_{|\mathcal{U}^r|} = \mathcal{U}^r$. Since $\mathcal{U}^r$ is locally formally of finite type this implies that $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}} = \bigcup_r \mathcal{U}^r$ is locally formally of finite type as well. This completes the proof of Theorem 4.13 □

Corollary 4.22. *The irreducible components of the topological space* $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}$ *are quasi-projective schemes over* ℓ*. In particular they are quasi-compact.*

Proof. Let T be an irreducible component and let x be its generic point. As in the proof of the theorem there is an r such that $x \in \mathcal{U}^r = \mathcal{U}_{r+d_0}^r \subset \mathcal{M}_{r+d_0}$. Since the underlying topological spaces of $\mathcal{M}_{r+d_0}$ and $\mathcal{M}^{r+d_0}$ coincide, are closed in $\underline{\mathcal{M}}_{\mathbb{L}_0}^{\hat{Z}}$, and quasi-projective over ℓ by Lemma 4.15, we see that $T \subset \mathcal{M}^{r+d_0}$ is a closed subscheme and the corollary follows. □

In the remainder of this section we fix an integer n and consider complete discrete valuation rings $\mathbb{F}_i[\![z_i]\!]$ for $i = 1, \ldots, n$ with finite residue fields $\mathbb{F}_i$, and fraction fields $Q_i = \mathbb{F}_i(\!(z_i)\!)$. Let $\mathbb{P}_i$ be a parahoric group scheme over $\operatorname{Spec} \mathbb{F}_i[\![z_i]\!]$, let $P_i := \mathbb{P}_i \times_{\mathbb{F}_i[\![z_i]\!]} \operatorname{Spec} \mathbb{F}_i(\!(z_i)\!)$, and let $\hat{Z}_i$ be a closed ind-subscheme of $\widehat{\mathcal{F}\ell}_{\mathbb{P}_i} := \mathcal{F}\ell_{\mathbb{P}_i} \widehat{\times}_{\mathbb{F}_i} \operatorname{Spf} \mathbb{F}_i[\![\zeta_i]\!]$ which is a bound in the sense of Definition 4.5. Let k be a field containing all $\mathbb{F}_i$. For all i let $\underline{\mathbb{L}}_i$ be a local $\mathbb{P}_i$-shtuka over k which is trivialized and decent. By Theorem 4.13 the Rapoport-Zink space $\mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i}$ is a formal scheme locally formally of finite type over $\operatorname{Spf} k[\![\zeta_i]\!]$. Therefore the product $\prod_i \mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i} := \mathcal{M}_{\underline{\mathbb{L}}_1}^{\hat{Z}_1} \widehat{\times}_k \ldots \widehat{\times}_k \mathcal{M}_{\underline{\mathbb{L}}_n}^{\hat{Z}_n}$ is a formal scheme locally formally of finite type over $\operatorname{Spf} k[\![\underline{\zeta}]\!] := \operatorname{Spf} k[\![\zeta_1, \ldots, \zeta_n]\!]$. Recall that the group $J_{\underline{\mathbb{L}}_i}(Q_i) = \operatorname{QIsog}_k(\underline{\mathbb{L}}_i)$ of quasi-isogenies of $\underline{\mathbb{L}}_i$ over k acts naturally on $\mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i}$; see Remark 4.14.

Let $\Gamma \subseteq \prod_i J_{\underline{\mathbb{L}}_i}(Q_i)$ be a subgroup which is discrete for the product of the z_i-adic topologies. We say that Γ is *separated* if it is separated in the profinite topology, that is, if for every $1 \ne g \in \Gamma$ there is a normal subgroup of finite index that does not contain g.

Proposition 4.23. *Let* $\Gamma \subseteq \prod_i J_{\underline{\mathbb{L}}_i}(Q_i)$ *be a separated discrete subgroup. Then the quotient* $\Gamma \backslash \prod_i \mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i}$ *is a locally noetherian, adic formal algebraic* $\operatorname{Spf} k[\![\underline{\zeta}]\!]$*-stack locally formally of finite type. Moreover, the 1-morphism* $\prod_i \mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i} \to \Gamma \backslash \prod_i \mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i}$ *is adic and étale.*

Here we say that a formal algebraic $\operatorname{Spf} k[\![\underline{\zeta}]\!]$-stack $\mathcal{X}$ (see [Har05, Definition A.5]) is *$\mathcal{J}$-adic* for a sheaf of ideals $\mathcal{J} \subset \mathcal{O}_{\mathcal{X}}$, if for some (any) presentation $X \to \mathcal{X}$ the formal scheme X is $\mathcal{J}\mathcal{O}_X$-adic, that is, $\mathcal{J}^r\mathcal{O}_X$ is an ideal of definition of X for all r. We then call $\mathcal{J}$ an *ideal of definition of $\mathcal{X}$*. We say that $\mathcal{X}$ is *locally formally of finite type* if $\mathcal{X}$ is locally noetherian, adic, and if the closed substack defined by the largest ideal of definition (see [Har05, A.7]) is an algebraic stack locally of finite type over $\operatorname{Spec} k$. Before proving the above proposition let us state the following lemma.

Lemma 4.24. *Let $\Gamma \subseteq \prod_i J_{\mathbb{L}_i}(Q_i)$ be a separated discrete subgroup. Let $U_i \subset \mathcal{F}\ell_{\mathbb{P}_i} \times_{\operatorname{Spec}\mathbb{F}_i} \operatorname{Spec} k$ be a quasi-compact subscheme and set $U = U_1 \times_k \ldots \times_k U_n$. Then the set*

$$\{\gamma \in \Gamma \colon \gamma U \cap U \neq \emptyset\}$$

is finite.

Proof. Note that $J_{\mathbb{L}_i}(Q_i)$ is contained in $LP_i(k^{\mathrm{alg}})$. By Theorem 4.4 any point $\underline{x} \in U(k^{\mathrm{alg}})$ can be represented by a tuple $(\underline{\mathcal{L}}_i, g_i)_i$, where $\underline{\mathcal{L}}_i$ is a trivialized local $\mathbb{P}_i$-shtuka over k^{alg} and the quasi-isogeny $g_i \colon \underline{\mathcal{L}}_i \to \underline{\mathbb{L}}_i$ is given by an element $g_i \in LP_i(k^{\mathrm{alg}})$. By [PR08, Theorem 1.4] the projection $LP_i \to \mathcal{F}\ell_{\mathbb{P}_i}$ admits sections locally for the étale topology, and hence étale locally on $\mathcal{F}\ell_{\mathbb{P}_i}$ the loop group LP_i is isomorphic to the product $\mathcal{F}\ell_{\mathbb{P}_i} \times_{\mathbb{F}_i} L^+\mathbb{P}_i$. In particular, the projection $\pi \colon \prod_i LP_i \to \prod_i \mathcal{F}\ell_{\mathbb{P}_i}$ is an affine morphism by [EGA, Proposition 2.7.1] and the preimage $\widetilde{U} \subseteq \prod_i LP_i$ of U under π is quasi-compact. Consider the morphism

$$\begin{array}{ccl} \widetilde{U} \times \widetilde{U} & \longrightarrow & \prod_i \mathcal{F}\ell_{\mathbb{P}_i} \\ (g_i\,, g_i')_i & \longmapsto & (g_i'\, g_i^{-1} \cdot L^+\mathbb{P}_i / L^+\mathbb{P}_i)_i . \end{array}$$

Since $\mathcal{F}\ell_{\mathbb{P}_i}$ is an ind-quasi-projective ind-scheme, this morphism factors through some quasi-projective subscheme $V \subset \prod_i \mathcal{F}\ell_{\mathbb{P}_i}$. Since $\underline{\mathbb{L}}_i$ is decent the group of quasi-isogenies $J_{\mathbb{L}_i}(Q_i) \subset LP_i(k^{\mathrm{alg}})$ is contained in $LP_i(\ell)$ for some finite field ℓ; see Remark 4.11. Let $\gamma \in \Gamma$ be such that $\underline{x} = (\underline{\mathcal{L}}_i, g_i)_i \in U$ and $\gamma\underline{x} = (\underline{\mathcal{L}}_i, \gamma_i g_i) \in U$ where $\gamma_i \in LP_i(\ell)$ is the projection of γ onto the i-th factor. Then $(g_i\,, \gamma_i g_i)_i \in \widetilde{U} \times \widetilde{U}$ and the image of γ under the projection map π lies in the finite set $V(\ell)$. Thus γ lies in the compact set $S = \pi^{-1}(V(\ell)) \cap \prod_i J_{\mathbb{L}_i}(Q_i)$. On the other hand Γ is discrete and thus has finite intersection with S. □

Proof of Proposition 4.23. By Lemma 4.21 there is a finite field ℓ_i and a constant d_i such that any ball in $\mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i}$ with radius d_i contains a rational point in $\mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i}(\ell_i)$. Let d be the maximum of the integers d_i and let ℓ be the compositum of the fields ℓ_i. Let $x := (x_i)_i$ be a point of $\prod_i \mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i}(\ell)$. We use the notation of the proof of Theorem 4.13 and consider in particular the metric $\tilde{d}_i$ on $\mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i}$ defined as in (4.5) after choosing some representation ι_i of $\mathbb{P}_i$. For a positive integer c we define the closed subscheme $B(x_i, c) := \{z \in \mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i} \text{ closed points} \colon \tilde{d}_i(z, x_i) \leq c\}$ of $(\mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i})_{\mathrm{red}}$. By Lemma 4.15 it is a quasi-projective scheme over $\operatorname{Spec}\ell$. In particular the set of ℓ-valued points $B(x_i, c)(\ell)$ is finite. Then for all $n \geq 2d$ the subscheme

$$\mathcal{U}_n^{2d}(x_i) \;:=\; B(x_i, n) \smallsetminus \bigcup_{y \in B(x_i, n+d)(\ell),\, \tilde{d}_i(y, x_i) > 2d} B(y, d) \;\subset\; (\mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i})_{\mathrm{red}}$$

is open in $B(x_i, n+d)$ and quasi-compact. Note that for $n \geq 3d$ all $\mathcal{U}_n^{2d}(x_i)$ coincide with $\mathcal{U}_{3d}^{2d}(x_i)$ by an argument similar to (4.6). Since $(\mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i})_{\mathrm{red}} = \varinjlim B(x_i, n+d)$ carries the limit topology, the subscheme $\mathcal{U}^{2d}(x_i) := \mathcal{U}_{3d}^{2d}(x_i) \subset (\mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i})_{\mathrm{red}}$ is open and quasi-compact. By Lemma 4.21 the union of the $\mathcal{U}^{2d}(x_i)$ for all $x_i \in \mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i}(\ell)$ equals $(\mathcal{M}_{\underline{\mathbb{L}}_i}^{\hat{Z}_i})_{\mathrm{red}}$.

For $x = (x_i)_i$ set $U_x := \prod_i \mathcal{U}^{2d}(x_i)$. Then $\gamma.U_x = U_{\gamma.x}$ and the open subsets U_x cover $\prod_i \mathcal{M}_{\mathbb{L}_i}^{\hat{Z}_i}$, for varying $x \in \prod_i \mathcal{M}_{\mathbb{L}_i}^{\hat{Z}_i}(\ell)$. Let $I \subset \prod_i \mathcal{M}_{\mathbb{L}_i}^{\hat{Z}_i}(\ell)$ be a set of representatives of the Γ-orbits in $\prod_i \mathcal{M}_{\mathbb{L}_i}^{\hat{Z}_i}(\ell)$. Fix an $x \in I$. Since Γ is separated, we may choose by Lemma 4.24 a normal subgroup $\Gamma'_x \subset \Gamma$ of finite index in Γ such that $U_x \cap \gamma' U_x = \emptyset$ for all $\gamma' \neq 1$ in Γ'_x. For all $\gamma \in \Gamma$ the natural morphism $\gamma U_x \to \Gamma'_x \backslash \prod_i \mathcal{M}_{\mathbb{L}_i}^{\hat{Z}_i}$ is an open immersion. Let V_x be the (finite) union of the images of these morphisms for all $\gamma \in \Gamma$. Then $(\Gamma'_x \backslash \Gamma) \backslash V_x$ is an open substack of $\Gamma \backslash \prod_i \mathcal{M}_{\mathbb{L}_i}^{\hat{Z}_i}$. Moreover, $(\Gamma'_x \backslash \Gamma) \backslash V_x$ is a finite étale quotient of V_x and the map $\coprod_{\gamma \in \Gamma} \gamma U_x \to (\Gamma'_x \backslash \Gamma) \backslash V_x$ is adic and étale. Therefore the morphism $\prod_i \mathcal{M}_{\mathbb{L}_i}^{\hat{Z}_i} \to \Gamma \backslash \prod_i \mathcal{M}_{\mathbb{L}_i}^{\hat{Z}_i}$ is adic and étale above $(\Gamma'_x \backslash \Gamma) \backslash V_x$. Since $(V_x)_{\mathrm{red}}$ is a scheme of finite type over k, its quotient $\big((\Gamma'_x \backslash \Gamma) \backslash V_x\big)_{\mathrm{red}} = (\Gamma'_x \backslash \Gamma) \backslash (V_x)_{\mathrm{red}}$ is an algebraic stack of finite type over k. Since the open subsets U_x cover $\prod_i \mathcal{M}_{\mathbb{L}_i}^{\hat{Z}_i}$ also the $(\Gamma'_x \backslash \Gamma) \backslash V_x$ cover $\Gamma \backslash \prod_i \mathcal{M}_{\mathbb{L}_i}^{\hat{Z}_i}$ and the proposition follows. □

5 The Relation Between Global $\mathfrak{G}$-Shtukas and Local $\mathbb{P}$-Shtukas

5.1 Preliminaries on $\mathfrak{G}$-Torsors

In this chapter we only assume that $\mathfrak{G}$ is a smooth affine group scheme over C. Let $\nu \in C$ be a closed point of C and set $C' := C \smallsetminus \{\nu\}$. We let $\mathscr{H}_e^1(C', \mathfrak{G})$ denote the category fibered in groupoids over the category of $\mathbb{F}_q$-schemes, such that $\mathscr{H}_e^1(C', \mathfrak{G})(S)$ is the full subcategory of $[C'_S/\mathfrak{G}](C'_S)$ consisting of those $\mathfrak{G}$-torsors over C'_S that can be extended to a $\mathfrak{G}$-torsor over the whole relative curve C_S. We denote by $\dot{(\ )}$ the restriction morphism

$$\dot{(\ )}\colon \mathscr{H}^1(C, \mathfrak{G}) \longrightarrow \mathscr{H}_e^1(C', \mathfrak{G})$$

which assigns to a $\mathfrak{G}$-torsor $\mathcal{G}$ over C_S the $\mathfrak{G}$-torsor $\dot{\mathcal{G}} := \mathcal{G} \times_{C_S} C'_S$ over C'_S. Let $\widetilde{\mathbb{P}}_\nu := \operatorname{Res}_{\mathbb{F}_\nu/\mathbb{F}_q} \mathbb{P}_\nu$ and $\widetilde{P}_\nu := \operatorname{Res}_{\mathbb{F}_\nu/\mathbb{F}_q} P_\nu$ be the Weil restrictions. Then $\widetilde{\mathbb{P}}_\nu$ is a smooth affine group scheme over $\operatorname{Spec} \mathbb{F}_q[\![z]\!]$. We apply Definition 2.2 for $\mathbb{F} = \mathbb{F}_q$ and let $\hat{\widetilde{\mathbb{P}}}_\nu := \widetilde{\mathbb{P}}_\nu \widehat{\times}_{\operatorname{Spec} \mathbb{F}_q[\![z]\!]} \operatorname{Spf} \mathbb{F}_q[\![z]\!] = \operatorname{Res}_{\mathbb{F}_\nu/\mathbb{F}_q} \hat{\mathbb{P}}_\nu$ be the ν-adic completion. We write $A_\nu \cong \mathbb{F}_\nu[\![z]\!]$ for a uniformizer $z \in \mathbb{F}_q(C)$. Then for every $\mathbb{F}_q$-algebra R we have

$$\begin{aligned} A_\nu \widehat{\otimes}_{\mathbb{F}_q} R &\cong (R \otimes_{\mathbb{F}_q} \mathbb{F}_\nu)[\![z]\!] &&= R[\![z]\!] \otimes_{\mathbb{F}_q} \mathbb{F}_\nu \quad \text{and} \\ Q_\nu \widehat{\otimes}_{\mathbb{F}_q} R &\cong (R \otimes_{\mathbb{F}_q} \mathbb{F}_\nu)(\!(z)\!) &&= R(\!(z)\!) \otimes_{\mathbb{F}_q} \mathbb{F}_\nu \,. \end{aligned}$$

This implies that

$$L^+\widetilde{\mathbb{P}}_\nu(R) = \widetilde{\mathbb{P}}_\nu(R[\![z]\!]) = \mathbb{P}_\nu(A_\nu \widehat{\otimes}_{\mathbb{F}_q} R) \quad \text{and} \quad L\widetilde{P}_\nu(R) = \widetilde{P}_\nu(R(\!(z)\!)) = P_\nu(Q_\nu \widehat{\otimes}_{\mathbb{F}_q} R).$$

If $\mathcal{G} \in \mathscr{H}^1(C, \mathfrak{G})(S)$, its completion $\widehat{\mathcal{G}}_\nu := \mathcal{G} \widehat{\times}_{C_S} (\operatorname{Spf} A_\nu \widehat{\times}_{\mathbb{F}_q} S)$ is a formal $\hat{\mathbb{P}}_\nu$-torsor (Definition 2.2) over $\operatorname{Spf} A_\nu \widehat{\times}_{\mathbb{F}_q} S$. The Weil restriction $\operatorname{Res}_{\mathbb{F}_\nu/\mathbb{F}_q} \widehat{\mathcal{G}}_\nu$ is a formal $\hat{\widetilde{\mathbb{P}}}_\nu$-torsor over $\operatorname{Spf} \mathbb{F}_q[\![z]\!] \widehat{\times}_{\mathbb{F}_q} S$ and corresponds by Proposition 2.4 to an $L^+\widetilde{\mathbb{P}}_\nu$-torsor over S which we denote $L_\nu^+(\mathcal{G})$. We obtain the morphism

$$L_\nu^+\colon \mathscr{H}^1(C, \mathfrak{G}) \longrightarrow \mathscr{H}^1(\operatorname{Spec} \mathbb{F}_q, L^+\widetilde{\mathbb{P}}_\nu)\,, \quad \mathcal{G} \mapsto L_\nu^+(\mathcal{G})\,.$$

Finally there is a morphism

$$L_\nu\colon \mathscr{H}_e^1(C', \mathfrak{G})(S) \longrightarrow \mathscr{H}^1(\operatorname{Spec} \mathbb{F}_q, L\widetilde{P}_\nu)(S)\,, \quad \dot{\mathcal{G}} \mapsto L_\nu(\dot{\mathcal{G}})$$

which sends the $\mathfrak{G}$-torsor $\dot{\mathcal{G}}$ over C'_S, having some extension $\mathcal{G}$ over C_S, to the $L\widetilde{P}_\nu$-torsor $L(L^+_\nu(\mathcal{G}))$ associated with $L^+_\nu\mathcal{G}$ under (2.1). Observe that $L_\nu(\mathcal{G})$ is independent of the extension $\mathcal{G}$. We therefore may write $L_\nu(\dot{\mathcal{G}}) := L(L^+_\nu(\mathcal{G}))$.

Lemma 5.1. *The above maps assign to each $\mathfrak{G}$-torsor $\mathcal{G}$ over C_S a triple $(\dot{\mathcal{G}}, L^+_\nu(\mathcal{G}), \varphi)$ where $\varphi\colon L_\nu(\dot{\mathcal{G}}) \xrightarrow{\sim} L(L^+_\nu(\mathcal{G}))$ is the canonical isomorphism of $L\widetilde{P}_\nu$-torsors. $\mathscr{H}^1(C,\mathfrak{G})(S)$ is equivalent to the category of such triples. In other words, the following diagram of groupoids is cartesian*

$$\begin{tikzcd}
\mathscr{H}^1(C,\mathfrak{G}) \arrow[r, "\dot{(\;)}"] \arrow[d, "L^+_\nu"'] & \mathscr{H}^1_e(C',\mathfrak{G}) \arrow[d, "L_\nu"] \\
\mathscr{H}^1(\operatorname{Spec}\mathbb{F}_q, L^+\widetilde{\mathbb{P}}_\nu) \arrow[r, "L"] & \mathscr{H}^1(\operatorname{Spec}\mathbb{F}_q, L\widetilde{P}_\nu)\,.
\end{tikzcd}$$

Proof. This follows from the glueing result of Beauville and Laszlo [BL95]. Indeed, we construct the inverse of the morphism $\mathscr{H}^1(C,\mathfrak{G}) \longrightarrow \mathscr{H}^1_e(C',\mathfrak{G}) \times_{\mathscr{H}^1(\operatorname{Spec}\mathbb{F}_q, L\widetilde{P}_\nu)} \mathscr{H}^1(\operatorname{Spec}\mathbb{F}_q, L^+\widetilde{\mathbb{P}}_\nu)$. Let S be an $\mathbb{F}_q$-scheme, $\dot{\mathcal{G}} \in \mathscr{H}^1_e(C',\mathfrak{G})(S)$, as well as $\mathcal{L}^+_\nu \in \mathscr{H}^1(\operatorname{Spec}\mathbb{F}_q, L^+\widetilde{\mathbb{P}}_\nu)(S)$ and $\varphi\colon L_\nu(\dot{\mathcal{G}}) \xrightarrow{\sim} L(\mathcal{L}^+_\nu)$ an isomorphism in $\mathscr{H}^1(\operatorname{Spec}\mathbb{F}_q, L\widetilde{P}_\nu)(S)$. Let $\mathcal{G}$ be an extension of $\dot{\mathcal{G}}$ to C_S. By Proposition 2.4 there is an étale covering $S' \to S$ and trivializations $\alpha\colon (\mathcal{L}^+_\nu)_{S'} \xrightarrow{\sim} (L^+\widetilde{\mathbb{P}}_\nu)_{S'}$ and $\beta\colon L^+_\nu(\mathcal{G})_{S'} \xrightarrow{\sim} (L^+\widetilde{\mathbb{P}}_\nu)_{S'}$ of the pullbacks to S'. We may assume that S' is the disjoint union of affine schemes of the form $\operatorname{Spec} R'$.

We glue $\dot{\mathcal{G}}_{S'}$ via the isomorphism $\alpha\circ\varphi\colon L_\nu(\dot{\mathcal{G}})_{S'} \xrightarrow{\sim} (L\widetilde{P}_\nu)_{S'}$ to $(L^+\widetilde{\mathbb{P}}_\nu)_{S'}$ as follows. Consider the algebraic torsor $\mathcal{G}\times_{C_S}\operatorname{Spec}(A_\nu\widehat{\otimes}_{\mathbb{F}_q}R')$ for the group scheme $\mathfrak{G}\times_C\operatorname{Spec}(A_\nu\widehat{\otimes}_{\mathbb{F}_q}R') = \mathbb{P}_\nu\times_{\operatorname{Spec}A_\nu}\operatorname{Spec}(A_\nu\widehat{\otimes}_{\mathbb{F}_q}R')$ over $\operatorname{Spec}(A_\nu\widehat{\otimes}_{\mathbb{F}_q}R')$. By [EGA, Proposition 2.7.1] it is affine of the form $\operatorname{Spec}B$. Its ν-adic completion $(\widehat{\mathcal{G}}_\nu)_{R'} = \mathcal{G}\,\widehat{\times}_{C_S}\operatorname{Spf}(A_\nu\widehat{\otimes}_{\mathbb{F}_q}R') = \operatorname{Spf}\widehat{B}$ is affine with $\widehat{B} = \varprojlim B/\nu^m B$. Recall that $L^+_\nu(\mathcal{G})_{R'}$ is the $L^+\widetilde{\mathbb{P}}_\nu$-torsor associated with the Weil restriction $\operatorname{Res}_{\mathbb{F}_\nu/\mathbb{F}_q}(\widehat{\mathcal{G}}_\nu)_{R'}$ by Proposition 2.4. The trivialization β induces a trivialization $\beta\colon \operatorname{Res}_{\mathbb{F}_\nu/\mathbb{F}_q}(\widehat{\mathcal{G}}_\nu)_{R'} \xrightarrow{\sim} \widehat{\widetilde{\mathbb{P}}}_\nu\,\widehat{\times}_{\operatorname{Spf}\mathbb{F}_q[\![z]\!]}\operatorname{Spf}R'[\![z]\!]$ which is determined by the section

$$\begin{aligned}
\beta^{-1}(1) &\in \operatorname{Hom}_{\operatorname{Spf}R'[\![z]\!]}(\operatorname{Spf}R'[\![z]\!]\,,\ \operatorname{Res}_{\mathbb{F}_\nu/\mathbb{F}_q}(\widehat{\mathcal{G}}_\nu)_{R'}) \\
&= \operatorname{Hom}_{\operatorname{Spf}(A_\nu\widehat{\otimes}_{\mathbb{F}_q}R')}(\operatorname{Spf}(A_\nu\widehat{\otimes}_{\mathbb{F}_q}R')\,,\ \operatorname{Spf}\widehat{B}) \\
&= \operatorname{Hom}^{\text{cont}}_{A_\nu\widehat{\otimes}_{\mathbb{F}_q}R'}(\widehat{B}\,,\ A_\nu\widehat{\otimes}_{\mathbb{F}_q}R') \\
&= \operatorname{Hom}_{A_\nu\widehat{\otimes}_{\mathbb{F}_q}R'}(B\,,\ A_\nu\widehat{\otimes}_{\mathbb{F}_q}R') \\
&= \operatorname{Hom}_{\operatorname{Spec}(A_\nu\widehat{\otimes}_{\mathbb{F}_q}R')}(\operatorname{Spec}(A_\nu\widehat{\otimes}_{\mathbb{F}_q}R')\,,\ \operatorname{Spec}B).
\end{aligned}$$

The latter induces a trivialization $\beta\colon \mathcal{G}\times_{C_S}\operatorname{Spec}(A_\nu\widehat{\otimes}_{\mathbb{F}_q}R') \xrightarrow{\sim} \mathbb{P}_\nu\times_{\operatorname{Spec}A_\nu}\operatorname{Spec}(A_\nu\widehat{\otimes}_{\mathbb{F}_q}R')$. Similarly the automorphism $\psi := \alpha\varphi\beta^{-1}$ of $(L\widetilde{P}_\nu)_{R'}$ is determined by the image

$$\psi(1) \;\in\; L\widetilde{P}_\nu(R') \;=\; P_\nu(\operatorname{Spec}Q_\nu\widehat{\otimes}_{\mathbb{F}_q}R')$$

and thus induces an automorphism ψ of $P_\nu\times_{\operatorname{Spec}Q_\nu}\operatorname{Spec}(Q_\nu\widehat{\otimes}_{\mathbb{F}_q}R')$. By [BL95] we may glue $\dot{\mathcal{G}}_{R'}$ with $\mathbb{P}_\nu\times_{\operatorname{Spec}A_\nu}\operatorname{Spec}(A_\nu\widehat{\otimes}_{\mathbb{F}_q}R')$ via the isomorphism $\psi\beta$ to obtain a uniquely determined $\mathfrak{G}$-torsor $\mathcal{G}'$ on $C_{R'}$.

We descend $\mathcal{G}'$ to C_S as follows. Let $S'' = S'\times_S S'$ and let $p_i\colon S''\to S'$ be the projection onto the i-th factor. Consider the element $h := (p_1^*\alpha\circ p_2^*\alpha^{-1})(1) \in L^+\widetilde{\mathbb{P}}_\nu(S'') = \mathbb{P}_\nu(A_\nu\widehat{\otimes}_{\mathbb{F}_q}\Gamma(S'',\mathcal{O}_{S''}))$. The isomorphism

$$(\,\mathrm{id}_{\dot{\mathcal{G}}_{S''}}, h\cdot)\colon p_2^*(\dot{\mathcal{G}}_{S'},\, \mathbb{P}_\nu\times_{\operatorname{Spec}A_\nu}\operatorname{Spec}(A_\nu\widehat{\otimes}_{\mathbb{F}_q}R'),\, \psi\beta) \xrightarrow{\sim} p_1^*(\dot{\mathcal{G}}_{S'},\, \mathbb{P}_\nu\times_{\operatorname{Spec}A_\nu}\operatorname{Spec}(A_\nu\widehat{\otimes}_{\mathbb{F}_q}R'),\, \psi\beta)$$

induces a descent datum on $\mathcal{G}'$ which is effective by [BLR90, §6.2, Theorem 7] because $\mathcal{G}'$ is affine over $C_{S'}$. Thus $\mathcal{G}'$ descends to a $\mathfrak{G}$-torsor $\mathcal{G} \in \mathscr{H}^1(C, \mathfrak{G})(S)$. This defines the inverse morphism and finishes the proof. $\square$

5.2 The Global-Local Functor

Analogously to the functor which assigns to an abelian variety A over a $\mathbb{Z}_p$-scheme its p-divisible group $A[p^\infty]$ we introduce a global-local functor from global $\mathfrak{G}$-shtukas to local $\mathbb{P}_{\nu_i}$-shtukas. But whereas abelian varieties only have one characteristic place, our global $\mathfrak{G}$-shtukas have n characteristic places $\underline{\nu} = (\nu_1, \ldots, \nu_n)$. So the global-local functor will assign to each global $\mathfrak{G}$-shtuka of characteristic $\underline{\nu}$ an n-tuple of local $\mathbb{P}_{\nu_i}$-shtukas. We begin with a

Remark 5.2. Let ν be a place on C and let $\mathbb{D}_\nu := \operatorname{Spec} A_\nu$ and $\hat{\mathbb{D}}_\nu := \operatorname{Spf} A_\nu$. Let $\deg \nu := [\mathbb{F}_\nu : \mathbb{F}_q]$ and fix an inclusion $\mathbb{F}_\nu \subset A_\nu$. Assume that we have a section $s\colon S \to C$ which factors through $\operatorname{Spf} A_\nu$, that is, the image in $\mathcal{O}_S$ of a uniformizer of A_ν is locally nilpotent. In this case we have

$$\hat{\mathbb{D}}_\nu \,\widehat{\times}_{\mathbb{F}_q} S \cong \coprod_{\ell \in \mathbb{Z}/(\deg \nu)} \mathrm{V}(\mathfrak{a}_{\nu,\ell}) \cong \coprod_{\ell \in \mathbb{Z}/(\deg \nu)} \hat{\mathbb{D}}_{\nu,S}, \tag{5.1}$$

where $\hat{\mathbb{D}}_{\nu,S} := \hat{\mathbb{D}}_\nu \,\widehat{\times}_{\mathbb{F}_\nu} S$ and where $\mathrm{V}(\mathfrak{a}_{\nu,\ell})$ denotes the component identified by the ideal $\mathfrak{a}_{\nu,\ell} = \langle a \otimes 1 - 1 \otimes s^*(a)^{q^\ell} : a \in \mathbb{F}_\nu \rangle$. Note that σ cyclically permutes these components and thus the $\mathbb{F}_\nu$-Frobenius $\sigma^{\deg \nu} =: \hat{\sigma}$ leaves each of the components $\mathrm{V}(\mathfrak{a}_{\nu,\ell})$ stable. Also note that there are canonical isomorphisms $\mathrm{V}(\mathfrak{a}_{\nu,\ell}) \cong \hat{\mathbb{D}}_{\nu,S}$ for all ℓ.

Although we will not need it in the sequel, we note the following interpretation of the component $\mathrm{V}(\mathfrak{a}_{\nu,0})$.

Lemma 5.3. *The section $s\colon S \to C$ induces an isomorphism of the component $\mathrm{V}(\mathfrak{a}_{\nu,0})$ with the formal completion $\widehat{C_S}^{\Gamma_s}$ of C_S along the graph Γ_s of s. In particular $\widehat{C_S}^{\Gamma_s}$ is canonically isomorphic to $\hat{\mathbb{D}}_{\nu,S}$.*

Proof. We first consider the formal completion $\widehat{C_{A_\nu}}^{\Gamma_{\tilde{s}}}$ of C_{A_ν} along the graph $\Gamma_{\tilde{s}}$ of the morphism $\tilde{s}\colon \operatorname{Spec} A_\nu \to C$. Let $\operatorname{Spec} A \subset C$ be a neighborhood of ν such that a uniformizing parameter z of C at ν lies in A. Then $I = (a \otimes 1 - 1 \otimes a : a \in A) \subset A \otimes_{\mathbb{F}_q} A_\nu$ is the ideal defining $\Gamma_{\tilde{s}}$ and $\widehat{C_{A_\nu}}^{\Gamma_{\tilde{s}}} = \operatorname{Spf} \varprojlim (A \otimes_{\mathbb{F}_q} A_\nu)/I^n$. The module I/I^2 is free of rank one over $A_\nu = (A \otimes_{\mathbb{F}_q} A_\nu)/I$ since C is a smooth curve over $\mathbb{F}_q$. We claim that I/I^2 is generated by $z \otimes 1 - 1 \otimes z$. Let $a \in A$, let $f \in \mathbb{F}_q(z)[X]$ be the minimal polynomial of a over $\mathbb{F}_q(z)$, and multiply it with the least common denominator to obtain the polynomial $F(X, z) \in \mathbb{F}_q[X, z]$. Note that the least common denominator lies in $\mathcal{O}_{C,\nu}^\times$ because a is integral over $\mathbb{F}_q[z]$ near ν. In $A \otimes_{\mathbb{F}_q} A_\nu[X]$ we use the abbreviations $\zeta := 1 \otimes z$ and $\alpha := 1 \otimes a$ and we consider the two-variable Taylor expansion of F at (α, ζ)

$$F(X, z \otimes 1) \;\equiv\; F(\alpha, \zeta) + \frac{\partial F}{\partial X}(\alpha, \zeta) \cdot (X - \alpha) + \frac{\partial F}{\partial z}(\alpha, \zeta) \cdot (z \otimes 1 - \zeta) \mod I^2$$

Plugging in $a \otimes 1$ for X yields $F(a \otimes 1, z \otimes 1) = 0$ in addition to $F(\alpha, \zeta) = 0$. Since A is unramified over $\mathbb{F}_q[z]$ at ν we have $\frac{\partial F}{\partial X}(\alpha, \zeta) \in A_\nu^\times$. This shows that $z \otimes 1 - 1 \otimes z$ generates the A_ν-module I/I^2.

By Nakayama's Lemma [Eis95, Corollary 4.7] there is an element $f \in 1 + I$ that annihilates the $A \otimes_{\mathbb{F}_q} A_\nu$-module $I/(z \otimes 1 - 1 \otimes z)$. We may replace $\operatorname{Spec}(A \otimes_{\mathbb{F}_q} A_\nu)$ by the principal open subset $\operatorname{Spec}(A \otimes_{\mathbb{F}_q} A_\nu)[\frac{1}{f}]$ which contains the graph $\Gamma_{\tilde{s}}$ and on which I is generated by the non-zero divisor $z \otimes 1 - 1 \otimes z$. This implies that also I^n/I^{n+1} is a free A_ν-module with generator $(z \otimes 1 - 1 \otimes z)^n$.

Therefore the morphism $A_\nu[\![t]\!] \to \varprojlim (A \otimes_{\mathbb{F}_q} A_\nu)/I^n, t \mapsto z \otimes 1 - 1 \otimes z$ induces an isomorphism on the associated graded rings, and hence is itself an isomorphism by [AM69, Lemma 10.23].

Now observe that $\mathrm{V}(\mathfrak{a}_{\nu,0})$ is the formal scheme on the topological space S with structure sheaf $\mathcal{O}_S[\![z]\!]$, and that s identifies the topological spaces S and Γ_s. Under base change to S this implies that the formal completion $\widehat{C_S}^{\Gamma_s}$ has structure sheaf $\mathcal{O}_S[\![t]\!] = \mathcal{O}_S[\![z-\zeta]\!]$, where we write $t = z - \zeta$. Since ζ is locally nilpotent in $\mathcal{O}_S$, the latter is isomorphic to $\mathcal{O}_S[\![z]\!]$ proving the lemma. ☐

Definition 5.4. Fix a tuple $\underline{\nu} := (\nu_i)_{i=1\ldots n}$ of places on C with $\nu_i \neq \nu_j$ for $i \neq j$. Let $A_{\underline{\nu}}$ be the completion of the local ring $\mathcal{O}_{C^n,\underline{\nu}}$ of C^n at the closed point $\underline{\nu}$, and let $\mathbb{F}_{\underline{\nu}}$ be the residue field of the point $\underline{\nu}$. Then $\mathbb{F}_{\underline{\nu}}$ is the compositum of the fields $\mathbb{F}_{\nu_i}$ inside $\mathbb{F}_q^{\mathrm{alg}}$, and $A_{\underline{\nu}} \cong \mathbb{F}_{\underline{\nu}}[\![\zeta_1, \ldots, \zeta_n]\!]$ where ζ_i is a uniformizing parameter of C at ν_i. Let the stack

$$\nabla_n \mathscr{H}^1(C, \mathfrak{G})^{\underline{\nu}} \;:=\; \nabla_n \mathscr{H}^1(C, \mathfrak{G}) \,\widehat{\times}_{C^n} \operatorname{Spf} A_{\underline{\nu}}$$

be the formal completion of the ind-algebraic stack $\nabla_n \mathscr{H}^1(C, \mathfrak{G})$ along $\underline{\nu} \in C^n$. It is an ind-algebraic stack over $\operatorname{Spf} A_{\underline{\nu}}$ which is ind-separated and locally of ind-finite type by Theorem 2.15. Set $\mathbb{P}_{\nu_i} := \mathfrak{G} \times_C \operatorname{Spec} A_{\nu_i}$ and $\hat{\mathbb{P}}_{\nu_i} := \mathfrak{G} \times_C \operatorname{Spf} A_{\nu_i}$.

Let $(\mathcal{G}, s_1, \ldots, s_n, \tau) \in \nabla_n \mathscr{H}^1(C, \mathfrak{G})^{\underline{\nu}}(S)$, that is, $s_i \colon S \to C$ factors through $\operatorname{Spf} A_{\nu_i}$. By Remark 5.2 we may decompose $\mathcal{G} \,\widehat{\times}_{C_S} (\operatorname{Spf} A_{\nu_i} \,\widehat{\times}_{\mathbb{F}_q} S) \cong \coprod_{\ell \in \mathbb{Z}/(\deg \nu_i)} \mathcal{G} \,\widehat{\times}_{C_S} \mathrm{V}(\mathfrak{a}_{\nu_i,\ell})$ with components $\mathcal{G} \,\widehat{\times}_{C_S} \mathrm{V}(\mathfrak{a}_{\nu_i,\ell}) \in \mathscr{H}^1(\hat{\mathbb{D}}_{\nu_i}, \hat{\mathbb{P}}_{\nu_i})$. Using Proposition 2.4 we view $\big(\mathcal{G} \,\widehat{\times}_{C_S} \mathrm{V}(\mathfrak{a}_{\nu_i,0}), \tau^{\deg \nu_i}\big)$ as a local $\mathbb{P}_{\nu_i}$-shtuka over S, where $\tau^{\deg \nu_i} \colon (\sigma^{\deg \nu_i})^* \mathcal{L}_i \xrightarrow{\sim} \mathcal{L}_i$ is the $\mathbb{F}_{\nu_i}$-Frobenius on the loop group torsor $\mathcal{L}_i$ associated with $\mathcal{G} \,\widehat{\times}_{C_S} \mathrm{V}(\mathfrak{a}_{\nu_i,0})$. We define the *global-local functor* $\widehat{\underline{\Gamma}}$ by

$$\begin{aligned}
&\widehat{\Gamma}_{\nu_i} \colon \nabla_n \mathscr{H}^1(C, G)^{\underline{\nu}}(S) \longrightarrow Sht_{\mathbb{P}_{\nu_i}}^{\operatorname{Spec} A_{\nu_i}}(S), \quad (\mathcal{G}, \tau) \longmapsto \big(\mathcal{G} \,\widehat{\times}_{C_S} \mathrm{V}(\mathfrak{a}_{\nu_i,0}), \tau^{\deg \nu_i}\big) \quad \text{and} \\
&\widehat{\underline{\Gamma}} \;:=\; \prod_i \widehat{\Gamma}_{\nu_i} \colon \nabla_n \mathscr{H}^1(C, G)^{\underline{\nu}}(S) \longrightarrow \prod_i Sht_{\mathbb{P}_{\nu_i}}^{\operatorname{Spec} A_{\nu_i}}(S)\,. \qquad (5.2)
\end{aligned}$$

Note that $\widehat{\Gamma}_{\nu_i}$ and $\widehat{\underline{\Gamma}}$ also transform quasi-isogenies into quasi-isogenies.

Remark 5.5. Consider the preimages in $\mathrm{V}(\mathfrak{a}_{\nu_i,\ell})$ of the graphs $\Gamma_{s_j} \subset C_S$ of s_j. Since $\nu_i \neq \nu_j$ for $i \neq j$ the preimage of Γ_{s_j} is empty for $j \neq i$. Also the preimage of Γ_{s_i} is contained in $\mathrm{V}(\mathfrak{a}_{\nu_i,0})$ and does not meet $\mathrm{V}(\mathfrak{a}_{\nu_i,\ell})$ for $\ell \neq 0$. Thus for $\ell \neq 0$ the restriction of τ to $\mathrm{V}(\mathfrak{a}_{\nu_i,\ell})$ is an isomorphism

$$\tau \times \mathrm{id} \colon \sigma^*\big(\mathcal{G} \,\widehat{\times}_{C_S} \mathrm{V}(\mathfrak{a}_{\nu_i,\ell-1})\big) = (\sigma^* \mathcal{G}) \,\widehat{\times}_{C_S} \mathrm{V}(\mathfrak{a}_{\nu_i,\ell}) \xrightarrow{\sim} \mathcal{G} \,\widehat{\times}_{C_S} \mathrm{V}(\mathfrak{a}_{\nu_i,\ell})\,. \qquad (5.3)$$

This allows to recover $(\mathcal{G}, \tau) \,\widehat{\times}_{C_S} (\operatorname{Spf} A_{\nu_i} \,\widehat{\times}_{\mathbb{F}_q} S)$ from $\big(\mathcal{G} \,\widehat{\times}_{C_S} \mathrm{V}(\mathfrak{a}_{\nu_i,0}), \tau^{\deg \nu_i}\big)$ via the isomorphism

$$\prod_\ell (\tau^\ell \,\widehat{\times}\, \mathrm{id}) \colon \left(\prod_\ell \sigma^{\ell *}\big(\mathcal{G} \,\widehat{\times}_{C_S} \mathrm{V}(\mathfrak{a}_{\nu_i,0})\big), \begin{pmatrix} 0 & & \tau^{\deg \nu_i} \\ 1 & \ddots & \\ & \ddots & \\ & 1 & 0 \end{pmatrix} \right) \xrightarrow{\sim} (\mathcal{G}, \tau) \,\widehat{\times}_{C_S} (\operatorname{Spf} A_{\nu_i} \,\widehat{\times}_{\mathbb{F}_q} S). \qquad (5.4)$$

Recall from Section 5.1 that the Weil restriction $\operatorname{Res}_{\mathbb{F}_{\nu_i}/\mathbb{F}_q} \widehat{\mathcal{G}}_\nu$ of the torsor $\widehat{\mathcal{G}}_\nu := \mathcal{G} \,\widehat{\times}_{C_S} (\operatorname{Spf} A_{\nu_i} \,\widehat{\times}_{\mathbb{F}_q} S)$ corresponds by Proposition 2.4 to an $L^+ \widetilde{\mathbb{P}}_{\nu_i}$-torsor $L^+_{\nu_i}(\mathcal{G})$. Then $(L^+_{\nu_i}(\mathcal{G}), \tau \,\widehat{\times}\, \mathrm{id})$ is a local $\widetilde{\mathbb{P}}_{\nu_i}$-shtuka over S. We call it the *local $\widetilde{\mathbb{P}}_{\nu_i}$-shtuka associated with $\underline{\mathcal{G}}$ at the place ν_i*. By equation (5.4) there is an equivalence between the category of local $\mathbb{P}_{\nu_i}$-shtukas over schemes $S \in \mathcal{N}ilp_{A_{\nu_i}}$ and the category of local $\widetilde{\mathbb{P}}_{\nu_i}$-shtukas over S for which the Frobenius τ is an isomorphism outside $\mathrm{V}(\mathfrak{a}_{\nu_i,0})$. (Compare also [BH11, Proposition 8.8].)

Remark 5.6. Note that in a similar way one can associate a local $\widetilde{\mathbb{P}}_\nu$-shtuka $L^+_\nu(\underline{\mathcal{G}})$ with a global $\mathfrak{G}$-shtuka $\underline{\mathcal{G}} = (\mathcal{G}, \tau)$ at a place ν outside the characteristic places ν_i. Namely $L^+_\nu(\underline{\mathcal{G}})$ is the local $\widetilde{\mathbb{P}}_\nu$-shtuka associated with $\operatorname{Res}_{\mathbb{F}_\nu/\mathbb{F}_q}\big(\underline{\mathcal{G}}\,\widehat{\times}_{C_S}(\operatorname{Spf} A_\nu\,\widehat{\times}_{\mathbb{F}_q} S)\big)$ by Proposition 2.4. It is étale because τ is an isomorphism at ν. We call $L^+_\nu(\underline{\mathcal{G}})$ the *étale local* $\widetilde{\mathbb{P}}_\nu$*-shtuka associated with* $\underline{\mathcal{G}}$ *at the place* $\nu \notin \underline{\nu}$. In [AH13, Chapter 6] it will become useful for considerations of Tate-modules (Definition 3.5).

For this purpose we write $A_\nu \cong \mathbb{F}_\nu[\![z]\!]$. For every representation $\rho\colon \mathbb{P}_\nu \to \mathrm{GL}_{r,A_\nu}$ in $\operatorname{Rep}_{A_\nu}\mathbb{P}_\nu$ we consider the representation $\tilde\rho \in \operatorname{Rep}_{\mathbb{F}_q[\![z]\!]}\widetilde{\mathbb{P}}_\nu$ which is the composition of $\operatorname{Res}_{\mathbb{F}_\nu/\mathbb{F}_q}(\rho)\colon \widetilde{\mathbb{P}}_\nu \to \operatorname{Res}_{\mathbb{F}_\nu/\mathbb{F}_q}\mathrm{GL}_{r,A_\nu}$ followed by the natural inclusion $\operatorname{Res}_{\mathbb{F}_\nu/\mathbb{F}_q}\mathrm{GL}_{r,A_\nu} \subset \mathrm{GL}_{r\cdot[\mathbb{F}_\nu:\mathbb{F}_q]\,,\,\mathbb{F}_q[\![z]\!]}$. We set $\underline{\tilde{\mathcal{L}}} = L^+_\nu(\underline{\mathcal{G}})$ and $\check{\mathcal{T}}_{\underline{\tilde{\mathcal{L}}}}(\rho) := \check{\mathcal{T}}_{\underline{\tilde{\mathcal{L}}}}(\tilde\rho) := \check{T}_{\tilde\rho_*\underline{\tilde{\mathcal{L}}}}$. Then there is a canonical isomorphism $\check{\mathcal{T}}_{\underline{\tilde{\mathcal{L}}}}(\rho) \cong \varprojlim_n \rho_*\big(\underline{\mathcal{G}} \times_C \operatorname{Spec} A_\nu/(\nu^n)\big)^\tau$ of A_ν-modules.

If $\mathbb{F}_\nu \subset \mathcal{O}_S$ there also exists the decomposition (5.1) and we can associate a local $\mathbb{P}_\nu$-shtuka $\underline{\mathcal{L}}$ with $L^+_\nu(\underline{\mathcal{G}})$. The main difference to Definition 5.4 and Remark 5.5 is that there is no distinguished component of $\mathcal{G}\,\widehat{\times}_{C_S}(\operatorname{Spf} A_\nu\,\widehat{\times}_{\mathbb{F}_q} S)$, like the one given by the characteristic section at ν_i. But τ induces isomorphisms between all components as in (5.3). Therefore we may take any component and the associated local $\mathbb{P}_\nu$-shtuka $\underline{\mathcal{L}}$. Equation (5.4) shows that over $\mathbb{F}_\nu$-schemes S we obtain an equivalence between the category of étale local $\widetilde{\mathbb{P}}_\nu$-shtukas and the category of étale local $\mathbb{P}_\nu$-shtukas. There is also a canonical isomorphism of Tate functors $\check{\mathcal{T}}_{\underline{\mathcal{L}}} = \check{\mathcal{T}}_{L^+_\nu(\underline{\mathcal{G}})}$; compare [BH11, Proposition 8.5] for more details.

Like abelian varieties also global $\mathfrak{G}$-shtukas can be pulled back along quasi-isogenies of their associated local $\mathbb{P}$-shtukas.

Proposition 5.7. *Let* $\underline{\mathcal{G}} \in \nabla_n\mathscr{H}^1(C,\mathfrak{G})^{\underline{\nu}}(S)$ *be a global* $\mathfrak{G}$*-shtuka over* S *and let* $\nu \in C$ *be a place. Let* $L^+_\nu(\underline{\mathcal{G}})$ *be the local* $\widetilde{\mathbb{P}}_\nu$*-shtuka associated with* $\underline{\mathcal{G}}$ *at* ν *in the sense of Remark 5.5 (if* $\nu \in \underline{\nu}$*), respectively Remark 5.6 (if* $\nu \notin \underline{\nu}$*). Let* $\tilde f\colon \underline{\tilde{\mathcal{L}}}'_\nu \to L^+_\nu(\underline{\mathcal{G}})$ *be a quasi-isogeny of local* $\widetilde{\mathbb{P}}_\nu$*-shtukas over* S*. If* $\nu \in \underline{\nu}$ *we assume that the Frobenius of* $\underline{\tilde{\mathcal{L}}}'_\nu$ *is an isomorphism outside* $\mathrm{V}(\mathfrak{a}_{\nu,0})$*. If* $\nu \notin \underline{\nu}$ *we assume that* $\underline{\tilde{\mathcal{L}}}'_\nu$ *is étale. Then there exists a unique global* $\mathfrak{G}$*-shtuka* $\underline{\mathcal{G}}' \in \nabla_n\mathscr{H}^1(C,\mathfrak{G})^{\underline{\nu}}(S)$ *and a unique quasi-isogeny* $g\colon \underline{\mathcal{G}}' \to \underline{\mathcal{G}}$ *which is the identity outside* ν*, such that the local* $\widetilde{\mathbb{P}}_\nu$*-shtuka associated with* $\underline{\mathcal{G}}'$ *is* $\underline{\tilde{\mathcal{L}}}'_\nu$ *and the quasi-isogeny of local* $\widetilde{\mathbb{P}}_\nu$*-shtukas induced by* g *is* $\tilde f$*. We denote* $\underline{\mathcal{G}}'$ *by* $\tilde f^*\underline{\mathcal{G}}$*.*

Remark 5.8. Note that if $\nu \in \underline{\nu}$ then by Remark 5.5 there is an equivalence between the category of local $\widetilde{\mathbb{P}}_\nu$-shtukas over S for which the Frobenius τ is an isomorphism outside $\mathrm{V}(\mathfrak{a}_{\nu,0})$ and the category of local $\mathbb{P}_\nu$-shtukas over S. In particular, if $\widehat{\Gamma}_\nu(\underline{\mathcal{G}})$ is the local $\mathbb{P}_\nu$-shtuka associated with $\underline{\mathcal{G}}$ in Definition 5.4, then every isogeny $f\colon \underline{\mathcal{L}}'_\nu \to \widehat{\Gamma}_\nu(\underline{\mathcal{G}})$ corresponds under this equivalence to an isogeny $\tilde f\colon \underline{\tilde{\mathcal{L}}}'_\nu \to L^+_\nu(\underline{\mathcal{G}})$ as in the proposition. We obtain a global $\mathfrak{G}$-shtuka $\tilde f^*\underline{\mathcal{G}}$ which we also denote by $f^*\underline{\mathcal{G}}$. It satisfies $\widehat{\Gamma}_\nu(f^*\underline{\mathcal{G}}) = \underline{\mathcal{L}}'_\nu$ and $\widehat{\Gamma}_\nu(g\colon f^*\underline{\mathcal{G}} \to \underline{\mathcal{G}}) = f$.

Proof of Proposition 5.7. Let us set $\underline{\mathcal{G}} := (\mathcal{G}, \tau)$. Let $(\dot{\mathcal{G}}, L^+_\nu(\mathcal{G}), \varphi)$ be the triple for the place ν associated with the $\mathfrak{G}$-torsor $\mathcal{G}$ by Lemma 5.1. Thus $L^+_\nu(\underline{\mathcal{G}}) = (L^+_\nu(\mathcal{G}), \tau)$. We also set $\underline{\tilde{\mathcal{L}}}'_\nu = (\tilde{\mathcal{L}}'_\nu, \tau')$. Now the triple $(\dot{\mathcal{G}}, \tilde{\mathcal{L}}'_\nu, \tilde f^{-1}\varphi)$ defines a $\mathfrak{G}$-bundle $\mathcal{G}'$ over C_S which coincides with $\mathcal{G}$ over C'_S and inherits the Frobenius automorphism τ from $\underline{\mathcal{G}}$ over $C'_S \smallsetminus \bigcup_i \Gamma_{s_i}$. If $\nu \notin \underline{\nu}$ then this τ extends to an isomorphism over $\{\nu\} \times_{\mathbb{F}_q} S$ because $\underline{\tilde{\mathcal{L}}}'_\nu$ is étale. If $\nu \in \underline{\nu}$ then τ extends to an isomorphism over $C_S \smallsetminus \bigcup_i \Gamma_{s_i}$ because τ' is an isomorphism outside $\mathrm{V}(\mathfrak{a}_{\nu,0})$. This defines the global $\mathfrak{G}$-shtuka $\underline{\mathcal{G}}' \in \nabla_n\mathscr{H}^1(C,\mathfrak{G})^{\underline{\nu}}(S)$. The quasi-isogeny g is obtained from the identification $\dot{\mathcal{G}}' = \dot{\mathcal{G}}$. It has the desired properties. □

Finally we want to prove rigidity for quasi-isogenies of global $\mathfrak{G}$-shtukas. This is the global counterpart of Proposition 2.11 and fits into the analogy between abelian varieties and global $\mathfrak{G}$-shtukas. It only holds in fixed finite characteristics.

Proposition 5.9. (Rigidity of quasi-isogenies for global $\mathfrak{G}$-shtukas) *Let S be a scheme in $\mathcal{N}ilp_{A_{\underline{\nu}}}$ and let $j\colon \bar{S} \to S$ be a closed immersion defined by a sheaf of ideals $\mathcal{I}$ which is locally nilpotent. Let $\underline{\mathcal{G}} = (\mathcal{G}, \tau)$ and $\underline{\mathcal{G}}' = (\mathcal{G}', \tau')$ be two global $\mathfrak{G}$-shtukas over S. Then*

$$\mathrm{QIsog}_S(\underline{\mathcal{G}}, \underline{\mathcal{G}}') \longrightarrow \mathrm{QIsog}_{\bar{S}}(j^*\underline{\mathcal{G}}, j^*\underline{\mathcal{G}}'), \quad f \mapsto j^*f$$

is a bijection of sets. f is an isomorphism at a place $\nu \notin \underline{\nu}$ if and only if j^*f is an isomorphism at ν. However this need not be true for places $\nu \in \underline{\nu}$.

Proof. It suffices to treat the case where $\mathcal{I}^q = (0)$. In this case the morphism σ_S factors through $j\colon \bar{S} \to S$

$$\sigma_S = j \circ \sigma' \colon S \to \bar{S} \to S.$$

Since $C_S \smallsetminus \cup_i \Gamma_{s_i} \supset C_S \smallsetminus \underline{\nu} \times_{\mathbb{F}_q} S$, the morphism τ defines a quasi-isogeny $\tau\colon \sigma'^* j^* \underline{\mathcal{G}} = \sigma_S^* \underline{\mathcal{G}} \to \underline{\mathcal{G}}$ which is an isomorphism outside $\underline{\nu} \times_{\mathbb{F}_q} S$ and similarly for $\underline{\mathcal{G}}'$. We fix a finite closed subset $D \subset C$ which contains all ν_i and consider quasi-isogenies which are isomorphisms outside D. Then the proposition follows from the diagram

$$\begin{tikzcd}
\mathcal{G}|_{C_S \smallsetminus D_S} \arrow[r, "f", "\cong"'] & \mathcal{G}'|_{C_S \smallsetminus D_S} \\
\sigma_S^*\mathcal{G}|_{C_S \smallsetminus D_S} \arrow[u, "\tau", "\cong"'] \arrow[r, "\sigma'^*(j^*f)", "\cong"'] & \sigma_S^*\mathcal{G}'|_{C_S \smallsetminus D_S} . \arrow[u, "\tau'", "\cong"']
\end{tikzcd}$$

□

5.3 The Analog of the Serre-Tate Theorem

The Serre-Tate Theorem relates the deformation theory of an abelian variety in characteristic p with the deformation theory of the associated p-divisible group. In this section we introduce the analogous situation over function fields and prove the analogous theorem relating the deformation theory of a global $\mathfrak{G}$-shtuka to the deformation theory of the associated n-tuple of local $\mathbb{P}_{\nu_i}$-shtukas via the global-local functor.

Let S be in $\mathcal{N}ilp_{A_{\underline{\nu}}}$ and let $j\colon \bar{S} \to S$ be a closed subscheme defined by a locally nilpotent sheaf of ideals $\mathcal{I}$. Let $\bar{\underline{\mathcal{G}}}$ be a global $\mathfrak{G}$-shtuka in $\nabla_n\mathscr{H}^1(C, \mathfrak{G})^{\underline{\nu}}(\bar{S})$. The category $Defo_S(\bar{\underline{\mathcal{G}}})$ of lifts of $\bar{\underline{\mathcal{G}}}$ to S consists of all pairs $(\underline{\mathcal{G}}, \alpha\colon j^*\underline{\mathcal{G}} \xrightarrow{\sim} \bar{\underline{\mathcal{G}}})$ where $\underline{\mathcal{G}}$ belongs to $\nabla_n\mathscr{H}^1(C, \mathfrak{G})^{\underline{\nu}}(S)$, where α is an isomorphism of global $\mathfrak{G}$-shtukas over $\bar{S}$, and where morphisms are isomorphisms between the $\underline{\mathcal{G}}$'s that are compatible with the α's.

Similarly for a local $\mathbb{P}$-shtuka $\bar{\underline{\mathcal{L}}}$ in $Sht_{\mathbb{P}}^{\mathbb{D}}(\bar{S})$ we define the category of lifts $Defo_S(\bar{\underline{\mathcal{L}}})$ of $\bar{\underline{\mathcal{L}}}$ to S. Notice that according to the rigidity of quasi-isogenies (Propositions 5.9 and 2.11) all Hom-sets in these categories contain at most one element.

Theorem 5.10. *Let $\bar{\underline{\mathcal{G}}} := (\bar{\mathcal{G}}, \bar{\tau})$ be a global $\mathfrak{G}$-shtuka in $\nabla_n\mathscr{H}^1(C, \mathfrak{G})^{\underline{\nu}}(\bar{S})$. Let $(\bar{\underline{\mathcal{L}}}_i)_i = \widehat{\underline{\Gamma}}(\bar{\underline{\mathcal{G}}})$. Then the functor*

$$Defo_S(\bar{\underline{\mathcal{G}}}) \longrightarrow \prod_i Defo_S(\bar{\underline{\mathcal{L}}}_i)\,, \quad \big(\underline{\mathcal{G}}, \alpha\big) \longmapsto \big(\widehat{\underline{\Gamma}}(\underline{\mathcal{G}}), \widehat{\underline{\Gamma}}(\alpha)\big)$$

induced by the global-local functor, is an equivalence of categories.

Proof. We proceed by constructing the inverse of the above functor. It suffices to treat the case where $\mathcal{I}^q = (0)$. In this case the morphism σ_S factors through $j\colon \bar{S} \to S$

$$\sigma_S = j \circ \sigma' \colon S \to \bar{S} \to S.$$

Let $(\underline{\mathcal{L}}_i, \hat{\alpha}_i \colon j^*\underline{\mathcal{L}}_i \xrightarrow{\sim} \underline{\bar{\mathcal{L}}}_i)_i$ be an object of $\prod_i Defo_S(\underline{\bar{\mathcal{L}}}_i)$. We consider the global $\mathfrak{G}$-shtuka $\underline{\widetilde{\mathcal{G}}} := \sigma'^*\underline{\bar{\mathcal{G}}}$ over S. Since $C_S \smallsetminus \cup_i \Gamma_{s_i} \supset C_S \smallsetminus \underline{\nu} \times_{\mathbb{F}_q} S$, the morphism $\bar{\tau}$ defines a quasi-isogeny $\bar{\tau}\colon j^*\underline{\widetilde{\mathcal{G}}} \to \underline{\bar{\mathcal{G}}}$ which is an isomorphism outside the graphs of the characteristic sections

$$\begin{tikzcd}
\sigma_S^* j^* \tilde{\mathcal{G}} \arrow[r, equal] \arrow[d] & \left(\sigma_S^2\right)^* \bar{\mathcal{G}} \arrow[r, "\sigma_S^*\bar{\tau}"] \arrow[d, "\sigma_S^*\bar{\tau}"'] & \sigma_S^* \bar{\mathcal{G}} \arrow[d, "\bar{\tau}"'] \\
j^* \tilde{\mathcal{G}} \arrow[r, equal] & \sigma_S^* \bar{\mathcal{G}} \arrow[r, "\bar{\tau}"] & \bar{\mathcal{G}}.
\end{tikzcd}$$

We write $\underline{\widehat{\Gamma}}(\underline{\widetilde{\mathcal{G}}}) = (\underline{\widetilde{\mathcal{L}}}_i)_i$ and $\underline{\bar{\mathcal{L}}}_i = (\bar{\mathcal{L}}_i, \hat{\bar{\tau}}_i)$. We compose $\hat{\bar{\tau}}_i^{-1}$ with $\hat{\alpha}_i$ to obtain quasi-isogenies $\hat{\bar{\gamma}}_i := \hat{\bar{\tau}}_i^{-1} \circ \hat{\alpha}_i \colon j^*\underline{\mathcal{L}}_i \to j^*\underline{\widetilde{\mathcal{L}}}_i$. By rigidity of quasi-isogenies (Proposition 2.11) they lift to quasi-isogenies $\hat{\gamma}_i \colon \underline{\mathcal{L}}_i \to \underline{\widetilde{\mathcal{L}}}_i$ with $j^*\hat{\gamma}_i = \hat{\bar{\gamma}}_i$. We put $\underline{\mathcal{G}} := \hat{\gamma}_n^* \circ \ldots \circ \hat{\gamma}_1^* \underline{\widetilde{\mathcal{G}}}$ (see Remark 5.8) and recall that there is a quasi-isogeny $\gamma\colon \underline{\mathcal{G}} \to \underline{\tilde{\mathcal{G}}}$ of global $\mathfrak{G}$-shtukas with $\underline{\widehat{\Gamma}}(\gamma) = (\hat{\gamma}_i)_i$ which is an isomorphism outside $\underline{\nu}$, see Proposition 5.7. We may now define the functor

$$\prod_i Defo_S(\underline{\bar{\mathcal{L}}}_i) \to Defo(\underline{\bar{\mathcal{G}}})$$

by sending $(\underline{\mathcal{L}}_i, \hat{\alpha}_i \colon j^*\underline{\mathcal{L}}_i \to \underline{\bar{\mathcal{L}}}_i)_i$ to $(\underline{\mathcal{G}}, \bar{\tau} \circ j^*\gamma)$. The quasi-isogeny $\alpha := \bar{\tau} \circ j^*\gamma$ is an isomorphism outside the graphs of the s_i by construction, and also at these graphs because $\underline{\widehat{\Gamma}}(\alpha) = (\hat{\alpha}_i)_i$. It can easily be seen by the above construction that these functors are actually inverse to each other. □

References

[AH13] M.E. Arasteh Rad, U. Hartl: *Uniformizing the moduli stacks of global $\mathfrak{G}$-Shtukas*, preprint on arXiv:.

[AM69] M.F. Atiyah, I.G. Macdonald: *Introduction to commutative algebra*, Addison-Wesley Publishing Co., Reading, Massachusets 1969.

[BL95] A. Beauville and Y. Laszlo: *Un lemme de descente*, Comptes Rendus Acad. Sci. Paris, vol. **320**, série I (1995), 335–340; also available at http://math1.unice.fr/∼beauvill/bibli.html.

[BD] A. Beilinson, V. Drinfeld: *Quantization of Hitchin's integrable system and Hecke eigensheaves*, preprint at http://www.math.uchicago.edu/∼mitya/langlands.html.

[BH11] M. Bornhofen, U. Hartl: *Pure Anderson motives and abelian τ-sheaves*, Math. Z. **268** (2011), 67–100; also available as arXiv:0709.2809.

[BLR90] S. Bosch, W. Lütkebohmert, M. Raynaud: *Néron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **21**, Springer-Verlag, Berlin, 1990.

[BT72] F. Bruhat, J. Tits: *Groupes réductifs sur un corps local*, Inst. Hautes Études Sci. Publ. Math. **41** (1972), 5–251.

[Dri87] V. G. Drinfeld, *Moduli varieties of F-sheaves*, Func. Anal. and Appl. **21** (1987), 107–122.

[Eis95] D. Eisenbud: *Commutative Algebra with a View Toward Algebraic Geometry*, GTM **150**, Springer-Verlag, Berlin etc. 1995.

[Fal03] G. Faltings: *Algebraic loop groups and moduli spaces of bundles*, J. Eur. Math. Soc. **5** (2003), no. 1, 41–68.

[EGA] A. Grothendieck: *Élements de Géométrie Algébrique*, Publ. Math. IHES **4**, **8**, **11**, **17**, **20**, **24**, **28**, **32**, Bures-Sur-Yvette, 1960–1967; see also Grundlehren **166**, Springer-Verlag, Berlin etc. 1971.

[SGA 1] A. Grothendieck: *Revêtements étales et groupe fondamental*, LNM **224**, Springer-Verlag, Berlin-Heidelberg 1971; also available as arXiv:math/0206203.

[Hab78] W.J. Haboush: *Homogeneous vector bundles and reductive subgroups of reductive algebraic groups*, Amer. J. Math. **100** (1978), no. 6, 1123–1137.

[HR03] T. Haines and M. Rapoport: *On parahoric subgroups*, appendix to [PR08]; also available as arXiv:0804.3788.

[Har05] U. Hartl: *Uniformizing the Stacks of Abelian Sheaves*, in Number Fields and Function fields - Two Parallel Worlds, Papers from the 4th Conference held on Texel Island, April 2004, Progress in Math. **239**, Birkhäuser-Verlag, Basel 2005, pp. 167–222; also available as arXiv:math.NT/0409341.

[Har11] U. Hartl: *Period Spaces for Hodge Structures in Equal Characteristic*, Annals of Math. **173**, n. 3 (2011), 1241–1358; also available as arXiv:math.NT/0511686.

[HV11] U. Hartl, E. Viehmann: *The Newton stratification on deformations of local G-shtukas*, J. reine angew. Math. (Crelle) **656** (2011), 87–129; also available as arXiv:0810.0821.

[HV12] U. Hartl, E. Viehmann: *Foliations in deformation spaces of local G-shtukas*, Advances in Mathematics **229** (2012), 54–78; also available as arXiv:1002.2387.

[Kat73] N. Katz: *p-adic properties of modular schemes and modular forms*, Modular functions of one variable, III (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), LNM **350**, Springer, Berlin, 1973, pp. 69–190.

[Kot85] R. E. Kottwitz: *Isocrystals with additional structure*, Compositio Math. **56** (1985), no. 2, 201–220.

[Laf02] L. Lafforgue, *Chtoucas de Drinfeld et correspondance de Langlands*, Invent. Math. **147** (2002), 1–241; also available at http://www.ihes.fr/∼lafforgue/math/.

[Lan00] E. Landvogt: *Some Functorial Properties of Bruhat-Tits Buildings*, J. reine angew. Math. (Crelle) **518** (2000), 213–241.

[Lan56] S. Lang: *Algebraic groups over finite fields*, Amer. J. Math. **78** (1956), 555–563.

[LRS93] G. Laumon, M. Rapoport, U. Stuhler: *$\mathscr{D}$-elliptic sheaves and the Langlands correspondence*, Invent. Math. **113** (1993), 217–338.

[MFK94] D. Mumford, J. Fogarty, F. Kirwan: *Geometric invariant theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **34**, Springer-Verlag, Berlin, 1994.

[NN08] B.C. Ngô, T. Ngô Dac: *Comptage de G-chtoucas: La partie régulière elliptique*, J. Inst. of Math. Jussieu **7** (2008), n. 1, 181–203; also available at http://www.math.uchicago.edu/∼ngo/.

[NP01] B.C. Ngô, P. Polo: *Résolutions de Demazure affines et formule de Casselman-Shalika géométrique*, J. Algebraic Geom. **10** (2001), no. 3, 515–547; also available at http://www.math.uchicago.edu/∼ngo/.

[PR08] G. Pappas, M. Rapoport: *Twisted loop groups and their affine flag varieties*, Advances in Math. **219** (2008), 118–198; also available as arXiv:math/0607130.

[PR09] R. Pink, E. Rütsche, *Image of the group ring of the Galois representation associated to Drinfeld modules*, J. Number Theory **129** (2009), no. 4, 866–881; also available at http://www.math.ethz.ch/∼pink/.

[PT06] R. Pink, M. Traulsen, *The Galois representations associated to a Drinfeld module in special characteristic, III. Image of the group ring*, J. Number Theory **116** (2006), no. 2, 373–395; also available at http://www.math.ethz.ch/∼pink/.

[RZ96] M. Rapoport, T. Zink: *Period Spaces for p-divisible Groups*, Ann. Math. Stud. **141**, Princeton University Press, Princeton 1996.

[RZ99] M. Rapoport, T. Zink: *A finiteness theorem in the Bruhat-Tits building: an application of Landvogt's embedding theorem*, Indag. Mathem. (N.S.) **10** (1999), 449–458.

[Ric10] T. Richarz: *Schubert varieties in twisted affine flag varieties and local models*, preprint on arXiv:1011.5416.

[Var04] Y. Varshavsky: *Moduli spaces of principal F-bundles*, Selecta Math. (N.S.) **10** (2004), no. 1, 131–166; also available as arXiv:math/0205130.

[Wed04] T. Wedhorn, *On Tannakian duality over valuation rings*, J. Algebra **282** (2004), no. 2, 575–609; also available at http://www2.math.uni-paderborn.de/people/torsten-wedhorn.

Esmail Mohammad Arasteh Rad
Universität Münster
Mathematisches Institut
Einsteinstr. 62
D – 48149 Münster
Germany

Urs Hartl
Universität Münster
Mathematisches Institut
Einsteinstr. 62
D – 48149 Münster
Germany
www.math.uni-muenster.de/u/urs.hartl/